

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

1. ...
 2. ...
 3. ...
 4. ...
 5. ...
 6. ...
 7. ...
 8. ...
 9. ...
 10. ...
 11. ...
 12. ...
 13. ...
 14. ...
 15. ...
 16. ...
 17. ...
 18. ...
 19. ...
 20. ...
 21. ...
 22. ...
 23. ...
 24. ...
 25. ...
 26. ...
 27. ...
 28. ...
 29. ...
 30. ...
 31. ...
 32. ...
 33. ...
 34. ...
 35. ...
 36. ...
 37. ...
 38. ...
 39. ...
 40. ...
 41. ...
 42. ...
 43. ...
 44. ...
 45. ...
 46. ...
 47. ...
 48. ...
 49. ...
 50. ...
 51. ...
 52. ...
 53. ...
 54. ...
 55. ...
 56. ...
 57. ...
 58. ...
 59. ...
 60. ...
 61. ...
 62. ...
 63. ...
 64. ...
 65. ...
 66. ...
 67. ...
 68. ...
 69. ...
 70. ...
 71. ...
 72. ...
 73. ...
 74. ...
 75. ...
 76. ...
 77. ...
 78. ...
 79. ...
 80. ...
 81. ...
 82. ...
 83. ...
 84. ...
 85. ...
 86. ...
 87. ...
 88. ...
 89. ...
 90. ...
 91. ...
 92. ...
 93. ...
 94. ...
 95. ...
 96. ...
 97. ...
 98. ...
 99. ...
 100. ...

1. ...
 2. ...
 3. ...
 4. ...
 5. ...
 6. ...
 7. ...
 8. ...
 9. ...
 10. ...
 11. ...
 12. ...
 13. ...
 14. ...
 15. ...
 16. ...
 17. ...
 18. ...
 19. ...
 20. ...
 21. ...
 22. ...
 23. ...
 24. ...
 25. ...
 26. ...
 27. ...
 28. ...
 29. ...
 30. ...
 31. ...
 32. ...
 33. ...
 34. ...
 35. ...
 36. ...
 37. ...
 38. ...
 39. ...
 40. ...
 41. ...
 42. ...
 43. ...
 44. ...
 45. ...
 46. ...
 47. ...
 48. ...
 49. ...
 50. ...
 51. ...
 52. ...
 53. ...
 54. ...
 55. ...
 56. ...
 57. ...
 58. ...
 59. ...
 60. ...
 61. ...
 62. ...
 63. ...
 64. ...
 65. ...
 66. ...
 67. ...
 68. ...
 69. ...
 70. ...
 71. ...
 72. ...
 73. ...
 74. ...
 75. ...
 76. ...
 77. ...
 78. ...
 79. ...
 80. ...
 81. ...
 82. ...
 83. ...
 84. ...
 85. ...
 86. ...
 87. ...
 88. ...
 89. ...
 90. ...
 91. ...
 92. ...
 93. ...
 94. ...
 95. ...
 96. ...
 97. ...
 98. ...
 99. ...
 100. ...

CONTRIBUTIONS TOWARD
A FLORA OF NEVADA. NO. 58.

- - -

ONAGRACEAE OF NEVADA

by

Philip A. Munz

November 1, 1941

Issued by

The Division of Plant Exploration and Introduction,
Bureau of Plant Industry,
U. S. Department of Agriculture,
Washington, D. C.

- - -

Work Projects Administration of Nevada,
Projects, O. P. 65-2-04-13, W. P. 658;
O. P. 165-2-04-21, W. P. 752.

- - -

Collaborator

University of Nevada

Address all queries concerning this publication to the Division
of Plant Exploration and Introduction, Bureau of Plant Industry,
U. S. Department of Agriculture, Washington, D. C.

CONTRIBUTIONS TOWARD
A FLORA OF NEVADA. NO. 32.

- - -

ONAGRACEAE OF NEVADA

by

Philip A. Munz

November 1, 1941

Issued by

The Division of Plant Exploration and Introduction,
Bureau of Plant Industry,
U. S. Department of Agriculture,
Washington, D. C.

- - -

Work Projects Administration of Nevada,
Projects, O. P. 65-2-04-13, W. P. 658;
O. P. 165-2-04-21, W. P. 752.

- - -

Collaborator

University of Nevada

Address all queries concerning this publication to the Division
of Plant Exploration and Introduction, Bureau of Plant Industry,
U. S. Department of Agriculture, Washington, D. C.

1911

1912

1913

1914

1915

1916

1917

1918

1919

ONAGRACEAE OF NEVADA

By Philip A. Munz*

Herbs or rarely shrubs, with simple alternate or opposite leaves; stipules none; flowers perfect, axillary or in terminal racemes, the parts mostly in 2's or in 4's; hypanthium adnate to ovary and usually prolonged beyond; sepals 4 (sometimes 2 or 5); petals 4 (sometimes 2 or 5), inserted at summit of hypanthium; stamens as many or twice as many as petals, borne at summit of hypanthium; ovary inferior, 4- (sometimes 2- or 5-) celled; style 1; stigma 4-lobed, or capitate, or discoid; fruit a capsule, rarely nut-like.

About 20 genera and perhaps 400 species of wide distribution, particularly well represented in western North America, including a number of plants of great use as ornamentals: Evening-primroses, Godetias, Fuchsias, etc.

KEY TO GENERA

- Flowers 2-merous; fruit indehiscent, obovoid, usually with hooked hairs9. Circaea.
- Flowers 4-merous.
- Seeds with tuft of hairs (coma) at one end.

* Professor of Botany, Pomona College, Claremont, California.

STATE OF MICHIGAN

IN SENATE,

January 11, 1906.

REPORT OF THE

COMMISSIONERS OF THE

STATE OF MICHIGAN,

FOR THE YEAR 1905.

ALBION: PRINTED BY THE STATE OF MICHIGAN, 1906.

THE STATE OF MICHIGAN,
COMMISSIONERS OF THE STATE OF MICHIGAN,
FOR THE YEAR 1905.

Hypanthium 2-3 cm. long and funnelform, with row of 8 scales within at about one-half its length; flowers scarlet.

1. Zauschneria.

Hypanthium less than 1 cm. long or lacking, without scales within; flowers not scarlet2. Epilobium.

Seeds without coma.

Fruit indehiscent, nut-like8. Gaura.

Fruit a dehiscent capsule.

Ovary 2-celled; hypanthium not prolonged beyond the ovary; flowers minute; stems with capillary branches7. Gayophytum.


Ovary 4-celled; hypanthium prolonged beyond the ovary. Anthers usually versatile, attached near the middle (difficult to determine in minute flowers); petals yellow or white, rarely red except on aging6. Oenothera.

Anthers innate, attached near the base, erect; petals white or pink. Sepals erect; petals small or wanting; pollen in tetrads3. Boisduvalia.

Sepals reflexed or the tips remaining united and turned to one side in anthesis; pollen not in tetrads.

Petals distinctly clawed, the claw at least one-fourth as long as the blade.

4. Clarkia.



Petals not at all or scarcely clawed, the
 claw not more than one-tenth as long
 as the blade. 5. Godetia.

1. ZAUSCHNERIA Presl.

ZAUSCHNERIA Presl, Rel. Haenk. 2: 28, pl. 52. 1831.

Erect or decumbent perennials, somewhat woody and with shredding bark at the base. Leaves sessile or nearly so, opposite or alternate, more or less fascicled. Inflorescence spicate, the flowers large, horizontal, fuchsia-like. Hypanthium scarlet, globose at base, then narrowed into a long tube bearing within the narrow part a transverse row of 8 lobe-like appendages, 4 erect and 4 deflexed. Sepals 4. Petals 4. Stamens 8, the alternate ones shorter; anthers versatile. Ovary 4-celled; stigma 4-lobed, peltate to capitate. Capsule imperfectly 4-celled, many-seeded. Seeds oblong, narrowed at base, comose at apex. (Named for Dr. M. Zauschner, a Professor of Natural History at the University of Prague).

Five species, one in Nevada.

1. ZAUSCHNERIA LATIFOLIA (Hook.) Greene, Pittonia 1: 25. 1887.

Z. californica var. latifolia Hook., Bot. Mag., pl. 4493.
 1850.

Z. californica subsp. latifolia (Hook.) Keck, Carnegie
 Inst. Pub. 520: 220. 1940.



Z. argentea A. Nels., Proc. Biol. Soc. Wash. 28: 173.

1905.


Stems 1-6 dm. high; plant green to grayish-green, villous to tomentose, often silky, often somewhat glandular; leaves mostly opposite, ovate to lance-ovate, tapering at both ends or rounded at base, 8-17 mm. wide, more or less denticulate with evident lateral veins; hypanthium scarlet, 2-3 cm. long; sepals erect, lanceolate, 8-10 mm. long; petals 2-cleft, 8-15 mm. long; capsule sessile or nearly so, linear, 4-angled, 8-nerved, with short beak, often curved, 1.5-2 cm. long, many-seeded; seeds broad, 1.5 mm. long.

Dry slopes in the Sierra Nevada of California. Apparently rare in Washoe County, Nevada.

2. EPILOBIUM L.

EPILOBIUM L. Sp. Pl. 347. 1753.

Usually herbaceous, sometimes suffruticose; annual, or more often perennial and wintering by rosettes, turions, etc. Leaves opposite or alternate, sessile or short-petioled, denticulate or entire. Flowers axillary or in terminal racemes or panicles, perfect. Hypanthium short or not prolonged beyond the ovary. Sepals 4. Petals 4, usually notched, purplish, pink, white or yellow. Stamens 8, the alternate ones shorter. Stigma oblong or 4-lobed. Capsule elongate, subcylindric to fusiform, or clavate, 4-celled, loculicidal. Seeds with tuft of silky hairs (coma) at upper end. (Greek, upon a pod; the flowers and capsule appearing together).



Over 100 species, cosmopolitan, except in the tropics.

KEY TO SPECIES

- 1. Hypanthium not prolonged above the ovary; flowers large, the petals 1-2 cm. long, entire, spreading.

(Subgenus Chamaenerion).

- 2. Style pilose at base, exceeding stamens; leaves 5-20 cm. long, membranaceous, reticulate-veiny beneath, with lateral veins confluent in marginal loops; racemes many-flowered, elongate, not leafy; seeds oblong, 1-1.3 mm. long.

1. E. angustifolium.

- 2. Style glabrous, shorter than stamens; leaves 2-6 cm. long, thick and fleshy, glaucous, not veiny; racemes few-flowered, short, leafy; seeds fusiform, 2 mm. long. .2. E. latifolium.

- 1. Hypanthium prolonged above the ovary; flowers usually smaller, the petals ascending. (Subgenus Euepilobium).

- 3. Flowers large, the petals 14-20 mm. long; stigma evidently lobed; leaves 1-2 cm. long, sessile, rounded at base.

3. E. obcordatum.


- 3. Flowers smaller, the petals 2-12 mm. long; stigma usually oblong.

- 4. Plant suffruticose, with several stems from a woody caudex, 1-3 dm. high; axils with fascicles of bract-like leaves.

4. E. nevadense.

- 4. Plant not suffruticose, or if slightly so, not with axillary fascicles.

5. Annuals; stems with exfoliating epidermis. Plants of dry situations.
6. Stems 3-9 dm. tall, glabrous except in upper parts; leaves usually alternate, with fascicles in axils; hypanthium 1-3 mm. long. 5. E. paniculatum.
6. Stems less than 3 dm. tall, puberulent throughout; leaves mostly opposite, without fascicles; hypanthium scarcely 1 mm. long. 6. E. minutum.
5. Perennials; epidermis not exfoliating. Mostly in moist situations.
7. Rootstocks bearing turions (globose or ovoid winter buds with fleshy overlapping scales) which may be rather loose in E. exaltatum.
8. Petals 5-9 mm. long; hypanthium about 3 mm. high, equally wide at summit; stems divaricately branched above. 7. E. exaltatum.
8. Petals shorter; hypanthium shorter, narrower.
9. Stems glabrous to pubescent, but not with decurrent lines of hair from leaf-bases; stems mostly simple. 8. E. brevistylum.
9. Stems with decurrent lines of hair from the leaf-bases.
10. Leaves linear to lance-linear, often with decurrent bases, not crowded, the margins often irregularly dentate. 9. E. Halleanum.
10. Leaves, at least the basal, ovate with rounded



- bases, often crowded, the leaf-margins mostly entire. 10. E. saximontanum.
7. Rootstocks not turioniferous, although sometimes with rather fleshy rosettes.
11. Plant pallid, glaucous and glabrous almost throughout. 11. E. glaberrimum.
11. Plant not glaucous, but green to canescent,
12. Stems mostly 1-2 dm. tall, simple above, with a few pairs of opposite leaves.
13. Leaves sessile, oblong or linear, suberect; stem slender. 12. E. oregonense.
13. Leaves more or less distinctly petioled and spreading.
14. Plant densely caespitose, stoloniferous; stems sigmoidally bent, mostly less than 1 dm. tall; petals purplish to rose, 4-6 mm. long; leaves 1-2 cm. long.
15. Capsule linear, slender, 1 mm. or less thick, 2-4 cm. long; seeds smooth, 1 mm. long; buds nodding. 13. E. alpinum.
15. Capsule subclavate, stouter, 1.5-2 mm. thick, 2-2.5 cm. long; seeds papillose, 1.5-2 mm. long; buds erect. . . 14. E. clavatum.
14. Plant not so densely caespitose; stems erect, 1-3 dm. tall; leaves 1.5-5 cm. long.
16. Petals purplish, 5-10 mm. long; seeds papil-

lose, 1 mm. long. . . . 15. E. Hornemannii.

16. Petals white or with pink tips, ca. 3 mm.

long; seeds smooth, 1 mm. long.

16. E. lactiflorum.

12. Stems mostly 3-10 dm. tall, usually freely branched, especially above; innovation by rosettes.

17. Inflorescence glandular-pubescent with spreading hairs. Common. 17. E. adenocaulon.

17. Inflorescence not glandular or only slightly so. Not common.

18. Floral branchlets, buds, sepals not usually white-pubescent; hairs usually spreading.

17a. E. adenocaulon var. perplexans.

18. Floral branchlets, etc. white-pubescent with incurved hairs. 18. E. californicum.


1. EPILOBIUM ANGUSTIFOLIUM L. Sp. Pl. 347. 1753.

Chamaenerion angustifolium Scop. Fl. Carn. ed. 2, 1: 271.

1772.

Epilobium spicatum Lam. Fl. Fr. 3: 482. 1778.

Perennial, with erect, mostly simple and few stems, 0.3-2.5 m. tall, glabrous below, commonly puberulent above; leaves alternate, lanceolate to linear-lanceolate, acute, nearly entire, sessile or nearly so, 5-15 (20) cm. long; flowers numerous in long terminal racemes, with small almost linear bracts; pedicels 5-12 mm. long; sepals lance-linear, 8-12 mm. long, commonly canescent-puberulent throughout, tinged lavender; petals li-



lac-purple, rose or even white, clawed, obovate, 8-18 mm. long; filaments dilated near base; stigma-lobes slender and elongate; capsule 5-7 cm. long; seeds oblong, 1-1.4 mm. long, with long dingy coma.

In fairly moist places, Transition to Hudsonian Zones.

Widely and generally distributed. North America, Eurasia.

Specimens seen from Ormsby, Storey, Douglas, Nye, Humboldt, Elko and Clark Counties, Nevada.

2. *EPILOBIUM LATIFOLIUM* L. Sp. Pl. 347. 1753.


Chamaenerion latifolium Sweet, Hort. Brit. ed. 2, 198. 1830.

Stems several from a caespitose rootstock, depressed or arched-ascending, 1-6 dm. tall, glabrous below, puberulent above; leaves elliptic-ovate to lanceolate, subopposite, fleshy, glaucous on both surfaces, entire, not veiny, acute, quite sessile, 2-6 cm. long; racemes short, few-flowered, leafy-bracted; pedicels 5-10 mm. long; sepals lanceolate, purplish, 13 mm. long; petals purple to rose to white, purple-veined, rhombic-ovate, 1.5-2.5 cm. long; style glabrous, shorter than stamens; stigma-lobes oblong; capsule canescent, 5-8 cm. long; seeds fusiform, 2 mm. long.

Wet places along streams, Arctic-Alpine Zone, North America and Eurasia. Seen from altitude of 10,000 feet, East Humboldt Mts., Elko Co., Nevada.

3. *EPILOBIUM OBCORDATUM* A. Gray, Proc. Amer. Acad. 6: 532. 1865.

Stems several from caespitose suffrutescent base, decumbent, 5-15 cm. tall, simple, glabrous below, usually puberulent above,



leafy; leaves opposite, usually crowded, glabrous and glaucous, ovate, almost entire, 1-2 cm. long, obtuse, sessile or nearly so; flowers 1 to few, born singly in uppermost axils; pedicels 2-20 mm. long, slender; hypanthium funnelform, 2-4 mm. long; sepals lanceolate, 9-12 mm. long; petals rose-purple, broadly obcordate, 12-20 mm. long; longer stamens two-thirds the petals; style equalling petals, glabrous, purplish; stigma-lobes short; capsule cylindrical-clavate, 2.5-3.5 cm. long; seeds 1.5 mm. long, finely papillose.

Ridges and slopes of Hudsonian and Arctic-Alpine Zones; central Oregon and Sierra Nevada of California, Nevada, and Idaho. Nevada collections seen from Washoe and Elko Counties.

4. *EPILOBIUM NEVADENSE* Munz, Bull. Torrey Bot. Club 56: 166.

1929.

Low, suffruticose, caespitose, from branched caudex, the branches prostrate, brown, with exfoliating bark, giving rise to stems 1-3 dm. long, leafy throughout; leaves subglabrous, denticulate, green or purplish, on short pubescent petioles; lower leaf-blades oblong, obtuse, opposite, 8-15 mm. long; principal cauline ones alternate, more narrow and more acute, with fascicles of bract-like leaves in axils; flowers few to several on each stem, in loose racemes, sessile or on short glandular pedicel-like stems; hypanthium glandular, 2-3 mm. long, tubular; sepals 3-5 mm. long; petals violet-purple, obcordate, 6-7 mm. long; capsules subfusiform, glandular-pubescent, 8-12 mm. long, 1.5-2 mm. thick; seeds smooth, brown, 1.5

mm. long.

Talus slopes, Upper Sonoran and Transition Zones, Charleston Mts., Clark Co., Nevada.

5. *EPILOBIUM PANICULATUM* Nutt. ex Torr. & Gray, Fl. No. Amer. 1:

490. 1840.

Annual, erect, the stem simple and with shreddy epidermis below, paniculately branched above, 3-9 (15) dm. tall, glabrous almost throughout; leaves linear-lanceolate to linear, 2-3 (5) cm. long, usually alternate, with fascicles of smaller ones in axils, short petioled, remotely denticulate; flowers in lax racemes on filiform branches; bracts subulate; pedicels 0.5-1.5 cm. long; glabrous; ovary glandular-puberulent; hypanthium 2-3 (4) mm. long; petals rose to lilac, 5-6 mm. long; style about $\frac{1}{2}$ the petals; capsule 2 cm. long, 4-angled, linear-clavate, beaked; seeds obovoid, flattened, 2 mm. long, with tawny coma.

A species of the western United States and British Columbia, growing on rather dry slopes and benches of the Upper Sonoran and Transition Zones. Known from Nevada in the following forms:

forma *ADENOCLADON* Haussknecht. Hypanthium 2-3 mm. long; pedicels and ovaries glandular-puberulent; petals colored. Material seen from Washoe, Humboldt, Lander, Nye, Eureka and Elko Counties.

forma *SUBULATUM* Haussknecht. Hypanthium 2-3 mm. long; pedicels glabrous; ovaries glabrous; petals colored. Nevada material seen from Washoe, Lander and Humboldt Counties.

forma TRACYI (Rydb.) St. John. Hypanthium 1-2 mm. long; pedicels and ovaries glabrous; petals white. Seen from Washoe Co.

forma LAEVICAULE (Rydb.) St. John. Hypanthium 4-5 mm. long; pedicels glabrous; ovaries glandular; petals rose. Seen from Elko Co.


6. *EPILOBIUM MINUTUM* Lindl. ex Hook. Fl. Bor. Am. 1: 207. 1834.

Annual, 5-30 cm. tall, usually diffusely branched, puberulent throughout; leaves mostly opposite, oblong-lanceolate to oblanceolate or lanceolate, entire or nearly so, 1-2 cm. long, on a much shorter petiole; flowers in axils of somewhat reduced upper leaves; pedicels 3-10 mm. long; hypanthium less than 1 mm. long; sepals about 1.5 mm. long; petals rose-lavender to white, emarginate, 2-4 mm. long; capsule subclavate, arcuate, 1.5-2.5 cm. long, beaked; seeds broadly obovoid, smooth, barely 1 mm. long.

British Columbia to central California, Montana and Arizona. To be expected from Nevada.

7. *EPILOBIUM EXALTATUM* Drew, Bull. Torrey Bot. Club 16: 151. 1889.

Perennial, with large turions; stems 3-9 dm. tall, rather slender, more or less pubescent, freely branched above with very slender branches; leaves lance-ovate, serrulate, nearly or quite sessile, 5-12 cm. long, the uppermost much reduced; flowers near ends of glandular-pubescent branches; pedicels 5-10 mm. long in fruit; hypanthium 2-3 mm. long, almost as



wide; sepals suberect, 3-4 mm. long; petals pink to rose-purple, 5-10 mm. long; capsules 3-5 cm. long; seeds beaked, rugose, 1 mm. long, coma white.

Wet places, Transition Zone, California to Washington and Idaho. Nevada material seen from Washoe, Ormsby, Esmeralda and Elko Counties.


8. *EPILOBIUM BREVISTYLUM* Barbey in Brewer & Watson, Bot. Calif. 1: 220. 1876.

Perennial, with compact turions, the dried scales of which persist at the base of the stem during the succeeding year; stems erect, simple or subsimple, slender, 2-6 dm. tall, glabrous below, crisp-pubescent or somewhat glandular about the inflorescence; leaves mostly opposite, ovate to elliptic-lanceolate or even linear-lanceolate, denticulate, with rounded sessile base, 2-4 cm. long, not crowded; flowers several; fruiting pedicels 5-15 mm. long; sepals 2-3 mm. long; petals purplish or paler, emarginate, 3-5 mm. long; capsules 4-6 cm. long; seeds about 1.5 mm. long, papillate.

Wet places, Transition to Hudsonian Zones, Washington to southern California, Montana and Colorado. Nevada material seen from Washoe, Ormsby, Lyon, Esmeralda, Humboldt and Elko Counties.

var. *URSINUM* (Parish) Jepson, Man. Fl. Pls. Calif., 670. 1925.

E. ursinum Parish ex Trelease, Ann. Rep. Mo. Bot. Garden 2: 100. 1891.



Lower stem and leaves pilose with spreading hairs. Range of the species. Nevada material seen from Washoe Co.

var. TENUE (Trel.) Jepson, l.c.

E. delicatum var. tenue Trelease, Ann. Rep. Mo. Bot. Gard.

2: 99. 1891.

Low, glabrous, 1-1.5 dm. tall; leaves oblong-linear, remote, subentire, erect, obtuse. Uncommon, with general range of the species. Nevada material seen from White Pine County.

9. EPILOBIUM HALLEANUM Hausskn., Mon. Epilobium. 261. 1884.

E. Drummondii Hausskn. l. c. 271.

Perennial with small turions; stems erect, simple, or nearly so, 1-4 dm. high, subglabrous below, with lines of hair from the decurrent bases of the leaves, glandular-puberulent in upper parts; leaves lance-linear, erect, some with clasping base, 1.5-4 cm. long, serrulate to entire, acute; flowers small; hypanthium 1-1.5 mm. long; sepals 2-3 mm. long; petals 2-4 mm. long, white to purplish; fruiting pedicels 3-5 mm. long; capsules 2-5 cm. long; seeds 1-1.5 mm. long, beaked.

Wet places in Transition Zone, British Columbia to California, Montana, Colorado. Collected by Watson in Diamond Vale, Elko Co.

10. EPILOBIUM SAXIMONTANUM Hausskn. Oesterr. Bot. Zeitschr. 29:

119. 1879.

E. ovatifolium Rydb. Bull. Torrey Bot. Club 31: 567. 1904.

1. The first part of the document discusses the importance of maintaining accurate records of all transactions and activities. It emphasizes that this is essential for transparency and accountability, particularly in the context of public institutions or organizations where trust is paramount. The text suggests that without proper record-keeping, it becomes difficult to track the flow of funds and resources, leading to potential mismanagement and loss of public confidence.

2. The second part of the document outlines the specific procedures and protocols that should be followed when conducting financial transactions. It details the roles and responsibilities of various stakeholders involved in the process, from the initiation of a request to the final approval and execution. The document stresses the need for strict adherence to these procedures to ensure consistency and fairness in all dealings.

3. The third part of the document addresses the issue of budgeting and financial planning. It discusses the importance of setting realistic goals and objectives based on available resources and market conditions. The text highlights that effective budgeting is not just about limiting expenses but also about maximizing the use of funds to achieve the organization's mission and vision.

4. The fourth part of the document focuses on the importance of regular audits and reviews. It explains that these processes are crucial for identifying areas of inefficiency, preventing fraud, and ensuring that the organization remains compliant with relevant laws and regulations. The document encourages a culture of continuous improvement and self-reflection to maintain high standards of performance.

5. The final part of the document provides a summary of the key points discussed and offers some concluding thoughts on the overall approach to financial management. It reiterates the message that success in any endeavor, especially one involving significant resources, depends on sound financial practices and a commitment to ethical conduct.

Perennial, turioniferous; stems simple or branched, often clustered, 1-4 dm. high, with decurrent lines of hair from leaf-bases; leaves opposite or upper ones alternate, the lower ones ovate, with rounded bases, upper may be elliptical or with subacute bases; leaves sessile or with petioles up to 2 mm. long, serrulate to entire, acute, 2-3 (5) cm. long, 1.2-1.5 cm. wide; petals white, pink or purplish, 3-5 mm. long; capsules 4-6 cm. long, crisp-hairy; seeds about 1 mm. long, brownish.

Wet places, Transition to Hudsonian Zones, Alberta to Colorado and Arizona. Material seen from Washoe, Elko and Clark Counties, Nevada.

11. *EPILOBIUM GLABERRIMUM* Barbey ex Brewer & Watson, Bot. Calif.

1: 220. 1876.

Perennial, with several stems from branching scaly root-stocks; stems simple or nearly so, slender, erect from somewhat decumbent base, glabrous and glaucous, sometimes slightly glandular-puberulent above, often purplish, 3-6 dm. tall; leaves pallid, glabrous, glaucous, ascending, oblong-lanceolate, obtuse, entire or nearly so, sessile, 2-5 cm. long, gradually reduced up the stem; flowers usually erect; sepals 1-2 mm. long; petals 4-7 mm. long, purplish to almost white; fruiting pedicels 1-2 cm. long; capsules 4-7 cm. long; seeds papillate, 1 mm. long.

Stream banks and wet places, Upper Transition to Hudsonian Zones, Washington to southern California, Idaho and Nevada. Seen from Washoe, Douglas, Ormsby and Storey Counties, Nevada.

var. FASTIGIATUM (Nutt.) Trel. Ann. Rep. Mo. Bot. Garden 2:

105. 1891.

E. affine var. fastigiatum Nutt. ex Torr. & Gray, Fl. No.

Amer. 1: 460. 1840.

E. platyphyllum Rydb., Bull. Torrey Bot. Club 40: 63. 1913.

Lower, 1-3 dm. tall; leaves broader, shorter, ovate, more crowded, 1.5-2.5 cm. long.

In similar situations to the species; British Columbia to Utah and California. Nevada material seen from Washoe and Elko Counties.

12. EPILOBIUM OREGONENSE Hausskn. Mon. Epilobium. 276. 1884.

Perennial, stoloniferous; stems simple, slender, erect, 1-3 dm. high, glabrous except for some sparse glandular pubescence in the inflorescence; leaves somewhat crowded on lower portion of stem, reduced and remote above, glabrous, oblong-linear to -ovate, entire to remotely denticulate, suberect, obtuse, sessile, 1-2.5 cm. long; flowers 1-few; fruiting pedicels 1-3.5 cm. long; sepals often purplish, 1-2 mm. long; petals cream to purplish, 4-7 mm. long, deeply emarginate; capsules erect, 2-5 cm. long, slender, often purplish; seeds smooth, blunt, scarcely 1 mm. long.

Boggy places, Upper Transition to Hudsonian Zones, British Columbia to southern California, Idaho and Nevada. Nevada material seen from Washoe and Esmeralda Counties.

13. EPILOBIUM ALPINUM L. Sp. Pl. 348. 1753.

E. anagallidifolium Lam. Dict. 2: 376. 1786.



Densely caespitose perennial, stoloniferous, with numerous simple, erect, slender, sigmoidally bent stems scarcely 1 dm. long and nodding at apex, glabrous or with pubescent lines; leaves divergent, rather uniformly distributed, oblong-ovate to -lanceolate, obtuse, entire or nearly so, 1-2 cm. long, on short petioles; inflorescence purplish, 1- to few-flowered, somewhat crisp-pubescent or even glandular; fruiting pedicels 5-15 mm. long; sepals 2 mm. long; petals lilac to purple, 4-5 mm. long; capsule slender, linear, 1 mm. thick, 2-4 cm. long; seeds smooth, 1 mm. long, with dingy coma.


Moist rockslides and stony places, Arctic-Alpine Zone, Alaska to central California, Colorado and Labrador. Eurasia. Seen from Washoe and Elko Counties, Nevada.

14. *EPILOBIUM CLAVATUM* Trel. Rep. Mo. Bot. Gard. 2: 111. pl. 48. 1891.

Habit much as in the preceding species, but stems 5-15 cm. tall, subglabrous to glandular-pubescent; leaves broadly ovate, obtuse, 1-2 cm. long; fruiting pedicels 1-2 cm. long; sepals 3-4 mm. long; petals purplish to rose-colored, 5-6 mm. long; capsule purplish, subclavate, 2-2.5 cm. long, stout, 1.5-2 mm. thick, frequently arcuate; seeds fusiform, papillose, 1.5-2 mm. long, with dingy coma.

Talus and slides, Arctic-Alpine Zone, British Columbia to Oregon, Montana and Colorado. Material seen from Elko Co., Nevada.

15. *EPILOBIUM HORNEMANNII* Reichenb. Ic. Crit. 2: 73. 1824.



Perennial with subterranean scaly branches; stems slender, erect, except at very base, simple, 1-3 dm. tall, glabrous except for the crisp pubescence on the decurrent lines, slightly glandular above; leaves ovate to elliptic-ovate, 1.5-4 cm. long, mostly obtuse, subentire to remotely denticulate, short-petioled; flowers erect; fruiting pedicels 1-2 cm. long; sepals 3-4 mm. long; petals rose or violet, 5-8 mm. long; capsules erect, linear, slender, less than 1 mm. thick, 4-5 cm. long; seeds 1 mm. long, papillose, with dingy coma.

Damp banks and meadows, Canadian and Hudsonian Zones, Alaska to central California, Colorado, New Hampshire, Greenland; Eurasia. Nevada material seen from Washoe, Ormsby, Elko and White Pine Cos.

16. *EPILOBIUM LACTIFLORUM* Hausskn. Oesterr. Bot. Zeitschr. 29: 89. 1879.

E. alpinum of American authors, not L.

Size and habit of the preceding species, but more glabrous on the decurrent lines as well as in the inflorescence; leaves delicate, elliptic or oblong-ovate, pale green, subentire or obscurely denticulate, obtuse, 2-5 cm. long; flowers few; petals 3 mm. long, white or rose-tipped; capsules slender, erect, linear, less than 1 mm. thick, 4-5 cm. long; seeds smooth, 1 mm. long with dingy coma.

Moist slopes and banks. Canadian to Arctic-Alpine Zones from Alaska to southern California, Colorado and New Hampshire; Eurasia. Seen from Washoe and Elko Cos., Nevada.

17. EPILOBIUM ADENOCAULON Hausskn. Oesterr. Bot. Zeitschr. 29: 119.
1879.

E. glandulosum var. adenocaulon Fernald, Rhodora 20: 35,
1918.

Perennial; stem erect, 3-10 dm. tall, glabrous below, glandular-pubescent and with few or no incurved hairs in inflorescence, simple or weakly branched below, freely branched above; innovations by rosettes; leaves glabrous or nearly so, ovate-to elliptic-lanceolate, 3-6 cm. long, obtuse to acute, serrulate, rounded into very short winged petioles, upper leaves gradually reduced and somewhat pubescent; sepals 2 mm. long; petals white to pale red, 4 mm. long; fruiting pedicels 3-8 mm. long; capsules slender, 4-6 cm. long, glabrate in age; seeds obovoid, 1 mm. long, abruptly short-beaked, with whitish coma.

Common in moist places, largely of the Transition Zone, British Columbia to southern California and Atlantic states. Seen from Washoe, Humboldt, Ormsby, Douglas, Storey, Lyon, Mineral, Nye, Churchill, Lander, Elko, White Pine and Clark Counties, Nevada.

var. OCCIDENTALE Trel. Ann. Rep. Mo. Bot. Gard. 2: 95. 1891.

E. occidentale Rydb., Mem. N.Y. Bot. Gard. 1: 275. 1900.

Leaves narrowly lanceolate, narrower than in the species; flowers purple or rose, the petals 5-6 mm. long.

British Columbia to central California and Utah. Seen from Esmeralda and Ormsby Cos., Nevada.



var. PERPLEXANS Trel. l.c. 96.

Usually less than 3 dm. tall, slender, rather simple, scarcely or not glandular; leaves thin, tapering at base to slender petiole; flowers whitish; petals 4 mm. long.

Transition Zone, eastern Washington and eastern California to Rocky Mts. Material from Nevada: Esmeralda, Lyon, Washoe, Mineral, Eureka and Elko Cos.

18. EPILOBIUM CALIFORNICUM Hausskn. Mon. Epilob. 260. 1884.

E. Palmeri Rydb. Bull. Torrey Bot. Club 31: 569. 1904.

With habit and stature of E. adenocaulon, but the upper parts strongly pubescent with white incurved non-glandular hairs; flowers tending to be smaller, less than 6 mm. long; petals white or pink, 2-4 mm. long.

Wet places of Upper Sonoran and Transition Zones, western Washington to California, mostly west of the Sierra Nevada, but apparently in Clark Co., Nevada.

3. BOISDUVALIA Spach

BOISDUVALIA Spach, Hist. Nat. Veg. Phan. 4: 383. 1835.

Caulescent, erect or spreading annuals; leaves alternate, simple, sessile; flowers small or minute in ours, in leafy spikes or axillary to foliage leaves; hypanthium produced above the ovary, short, funnelform; sepals 4, erect; petals 4, sessile, obovate, 2-lobed, purple to white; stamens 8, those opposite the petals shorter; anthers basifixed, all perfect; pollen in tetrads; stigma with 4 cuneate lobes; capsule 4-celled, 4-valved, sessile; seeds smooth,



without a coma. (Named for J. A. Boissieu, French naturalist and physician.)

Ten species, 3 in Nevada.

KEY TO SPECIES

1. Capsule septifragal, the septa wholly adherent to the placental axis, making the latter 4-winged; capsule not conspicuously beaked; floral bracts proportionally wider than cauline leaves. 1. B. densiflora.
1. Capsule loculicidal, the septa adhering to the valves at dehiscence; capsule beaked.
 2. Fruit straight, 5-8 mm. long; floral bracts ovate to oblong; petals 2-4 mm. long. 2. B. glabella.
 2. Fruit curved, 6-12 mm. long, the tip turned away from the stem; floral bracts linear; petals 1.5-2 mm. long.
 3. B. stricta.

1. BOISDUALIA DENSIFLORA (Lindl.) S. Wats. Bot. Calif. 1: 233.
1876.

Oenothera densiflora Lindl. Bot. Reg. 19: pl. 1593. 1833.

Simple or branched, particularly above, usually 3-10 dm. tall, villous, green to canescent, leafy throughout; lower leaves lanceolate to lance-linear, acute, entire, or denticulate, 2-5 cm. long, floral ones ovate, acute, 0.5-1.2 cm. long; inflorescence dense, long-spicate in fruit; sepals 2-4 mm. long; petals usually rose-purple, bilobed, 6-12 mm. long; capsule stout, straight, 8-10 mm. long, septifragal; seeds few,



angled, brown, paler at ends, concave on inner face, 1.5 mm. long.

In places that are moist in the early season, Upper Sonoran and Transition Zones, British Columbia to Lower California, Idaho and Nevada, from which state material has been seen from Washoe, Eureka and Elko Cos.

var. *IMBRICATA* Greene, Fl. Francisc., 225. 1891.

Floral bracts orbicular-ovate, so densely imbricated as quite to conceal the capsules.

With much the same range as the species. Nevada material seen from Storey Co.

var. *SALICINA* (Rydb.) Munz, Leaflets W. Bot. 3: 53. 1941.

B. salicina Rydb., Bull. Torrey Bot. Club 40: 62. 1913.

Plant canescent-strigose throughout, the pubescence short and not gland-tipped; petals mostly pale, rather short, 2.5-5 mm. long.

Ranging from Wash. to Calif., Idaho and Nevada. Nevada material seen from Washoe and Ormsby Cos.

2. *BOISDUVALIA GLABELLA* (Nutt.) Walp. Rep. 2: 89. 1843.

Oenothera glabella Nutt. ex Torr. & Gray, Fl. No. Amer. 1: 505.
1840.

Boisduvalia diffusa Greene, Proc. Acad. Philad. 1895: 547.
1896.

Simple or more frequently freely and decumbently branched

from base, 1-3 dm. tall, glabrous to pubescent, uniformly leafy; leaves sessile, lance-ovate to lance-oblong, acute, serrulate, bright green, 1-1.5 cm. long; flowers axillary, even in lowest axils; sepals 2 mm. long; petals purplish, 2-5 mm. long; capsule 6-8 mm. long, straight, pointed at tip; seeds numerous, grayish brown, narrowly subfusiform, angled, 1 mm. long.

Dry mud flats and vernal pools, Upper Sonoran Zone at scattered stations from British Columbia to southern California, Utah, North Dakota, Saskatchewan; Argentina. Nevada material seen from Washoe and Ormsby Cos.

3. *BOISDUVALIA STRICTA* (Gray) Greene, Fl. Francisc. 225. 1891.

Gayophytum strictum Gray, Proc. Amer. Acad. 7: 340. 1867.

Stems 1-4.5 dm. tall, simple or few-branched, pilose and quite canescent throughout; leaves linear to lanceolate, 1-4 cm. long, 2-3 mm. wide, acute, entire to sharply denticulate, nearly or quite sessile, the upper ones narrower than the lower ones; flowers axillary, often beginning near the base of the plant; sepals 1 mm. long; petals rose-purple to pale pink, 1.5-2 mm. long; capsule 8-10 mm. long, membranous, slender, usually curved outwards and attenuate, tardily loculicidal; seeds ovoid, brown, 1 mm. long, 6-8 in each cell.

In stream beds or other moist places which dry later in the season, Upper Sonoran and Transition Zones, Washington and Idaho to central California and Nevada. Seen from Humboldt and Elko Cos., Nevada.



4. CLARKIA Pursh

CLARKIA Pursh, Fl. Amer. Sept. 1: 260, pl. 11. 1814.

Annuals, simple or branched above, with spicate inflorescence and nodding or reflexed heads; hypanthium short or elongated; sepals 4, distinct or united in anthesis, petals unguiculate, the claws at least one-sixth the blades; blades simple or lobed, pink to purplish; stamens 4, alternate with the petals, or 8, with the epipetalous short and sometimes not functional; anthers basifixed; stigma 4-lobed; capsule linear or attenuate above, 4-celled; seeds in one row in each cell, cellular-pubescent and with reduced crest, or not pubescent and with minute transverse corrugations and conspicuous crest. (Named for Capt. William Clark of Lewis and Clark Expedition.)


Seven species, one in Nevada.

1. CLARKIA RHOMBOIDEA Dougl., in Hook. Fl. Bor. Am. 1: 214. 1834.

Godetia latifolia Nels. & Kennedy, Proc. Biol. Soc. Wash.

19: 156. 1906.

Two to 11 dm. tall, finely pubescent; leaves few, subopposite, lance-ovate to ovate-oblong or elliptic, the blades 2-7 cm. long, acute, entire, or remotely denticulate, subglabrous to finely pubescent; petioles 1-3 cm. long; hypanthium 1-3 mm. long, with scales and white hairs at summit; sepals green, usually distinct at anthesis; petals 5-10 mm. long, rose-purple, sometimes dotted, rhomboidal, with blade 2-4 times



as long as claw; stamens unequal, each with a scale at base; stigma-lobes rounded, 0.5 mm. long; capsules 1-3 cm. long, 2-4 mm. thick, 4-angled when dry; pedicels 1-4 mm. long; seeds brown, densely cellular-pubescent, 1 mm. long, with thickened ridge at summit and almost no cresting.

Rather dry slopes, Upper Sonoran and Transition Zones, Washington to Lower California, Arizona and South Dakota. Material seen from Washoe, Douglas, Ormsby and Humboldt Counties, Nevada.

5. *GODETIA* Spach

GODETIA Spach, Hist. Nat. Veg. Phan. 4: 386. 1835.

Annuals, mostly erect, with exfoliating epidermis on lower stems; leaves linear to spatulate, lower ones usually deciduous, upper ones reduced in size, secondary ones born in fascicles; inflorescence a spike or small panicle; flowers showy, white to rose to purple; hypanthium obconic to narrow-funneliform, with an inner ring of hair; sepals 4, distinct and reflexed in anthesis or wholly united and turned to one side; petals cuneate to obovate, entire to lobed, clawless or with short claw; stamens in 2 series, the opposite ones shorter; filaments filiform to flattened; anthers usually all fertile; capsule 4-sulcate, terete and 8-nerved, or heavily 8-ribbed, linear to ovoid, sessile to long pedicelled, beakless to long-beaked; seeds brown, sometimes cellular-puberulent, with fimbriate upper margin. (Named for C. H. Godet, 1797-1879, author of *Flora de Jura*.)

Genus of about 15 species, one in Nevada.

1. *GODETIA AMOENA* (Lehm.) G. Don in Sweet, Hort. Brit. ed. III,
237. 1839.

Oenothera amoena Lehm., Ind. Sem. Hort. Hamb. 8. 1821.

Erect, simple to diffusely branched, 1.5-10 dm. tall; leaf-blades lanceolate, 2-6 cm. long, 2-10 mm. wide, petioles 5-15 mm. long; buds erect or slightly drooped; hypanthium 4-10 mm. long, with inner ring of hairs one-third to one-half way from base; sepals 8-25 mm. long, generally united in anthesis; petals pink to purple, often with darker spot in center, cuneate to obovate, 1-4 cm. long; stigma lobes linear, yellow; capsule 1.5-4 cm. long, 2 mm. or more thick, linear, not enlarged above center, with or without short beak, usually pedicelled, deeply 4-sulcate when immature, terete and plainly nerved when mature or dry; seeds 1.5 mm. long, brown, with inconspicuous creasing.

Dry slopes, at edge of woods. Upper Sonoran and Transition Zones, British Columbia to central California and Nevada. Rare in Nevada; seen from Washoe and Ormsby Counties.

6. *OENOTHERA* L.

OENOTHERA L. Sp. Pl. 346. 1753.

Annual to perennial, caulescent or acaulescent herbs, with alternate or basal leaves; flowers in ours yellow to white, often aging red or purplish; hypanthium prolonged beyond the ovary, quite deciduous; sepals 4, reflexed in anthesis; petals 4; stamens 8,

equal, or if unequal, those opposite the petals shorter; anthers mostly versatile; stigma varying from 4 linear lobes to discoid or capitate; capsule membranous to woody, straight to curved or coiled, 4-celled, 4-valved, dehiscent; seeds many, naked. (Greek, wine-scenting, an ancient name given to some now unknown plant.)


Genus of perhaps 200 species, mostly of the temperate regions of the New World; 34 species in Nevada.

KEY TO SPECIES

1. Stigma with 4 linear lobes; flowers vespertine.
 2. Capsule broadly winged its entire length; seeds in 1 row in each cell, with corky tubercles; petals 4-5 cm. long.
(Subgenus *Megapterium*). 12. Oe. brachycarpa.
 2. Capsule not as above; seeds in 2 rows in each cell (except in *Anogra*), not with corky tubercles.
 3. Capsule appearing obpyramidal because of the winged angles in the upper part; seeds with wing-like margin at the summit; petals 1-2 cm. long. (Subgenus *Lavauxia*.)
11. Oe. flava.
 3. Capsule cylindrical to lance-ovoid, not winged; seeds not winged.
 4. Plants tufted or almost acaulescent; seeds with deep furrow along the raphe; capsule cylindrical or lance-ovoid, thick-walled, usually ridged, often tubercled.
(Subgenus *Pachylophis*.)
 5. Flowers white.



6. Petals 2.5-5 cm. long; hypanthium 5-8 cm. long; perennial. 7. Oe. caespitosa.
6. Petals 0.8-1.2 cm. long; hypanthium 2.5-4 cm. long; annual. 8. Oe. cavernae.
5. Flowers yellow.
7. Capsule 3.5-6 cm. long; plant with dense short soft pubescence; leaves with large terminal lobe and few small lateral ones. Transition Zone.
10. Oe. xylocarpa.
7. Capsule 2-3.5 cm. long; plant villous or pilose; leaves rather deeply and regularly pinnatifid. Lower Sonoran Zone. 9. Oe. primiveris.
4. Plants definitely caulescent; seeds not with deep raphal furrow.
8. Seeds sharply angled, horizontal; flowers yellow; capsule gradually tapering upward. (Subgenus *Eucenothera*.)
9. Petals 1-2 cm. long; sepals 1-1.5 cm. long; hypanthium 3-4 cm. long. 1. Oe. Rydbergii.
9. Petals 2.5-4 cm. long; sepals 2.5-5 cm. long.
10. Free tube of hypanthium 3-5 cm. long; leaves usually distinctly denticulate. 2. Oe. Hookeri.
10. Free tube of hypanthium 8-12 cm. long; leaves often almost entire. 3. Oe. longissima.
8. Seeds not sharply angled, ascending; flowers white;



capsule subcylindric; seeds in 1 row in each cell

(Subgenus Anogra.)

11. Spring or winter annuals, coarse; basal leaves rhombic, the blades 2-8 cm. long; capsules woody with exfoliating epidermis and 2-7 cm. long.

4. Oe. deltoides.

11. Perennials; basal leaves smaller and more narrow; capsules not woody.

12. Plants essentially glabrous; capsules frequently contorted, seeds linear-obovoid. 5. Oe. pallida.

12. Plants canescent to hoary or villous; capsules spreading, usually not contorted; seeds plump and obovoid. 6. Oe. californica.

1. Stigma capitate, discoid, or 4-toothed; flowers mostly diurnal; seeds in one row in each cell.

13. Ovary with a long narrow upper sterile part grading into the enlarged basal fertile part; plants quite acaulescent; flowers yellow. (Subgenus Taraxia.)

14. Annuals; capsules broadly and truncately 4-winged, not over 1 cm. long; petals 2.5-3 mm. long. . . . 14. Oe. Palmeri.

14. Perennials; capsules somewhat cylindrical, at most angled, not winged, usually more than 1 cm. long; petals 8-15 mm. long.

15. Leaves entire or with few teeth; capsule glabrous.

15. Oe. heterantha.

15. Leaves deeply pinnatifid; capsule densely pubescent.

16. Oe. tanacetifolia.

13. Ovary not with long upper sterile part.
16. Stigma discoid, somewhat shallowly 4-lobed; hypanthium 2.5-5 cm. long (Subgenus *Salpingia*). 13. Oe. lavandulaefolia.
16. Stigma capitate; hypanthium much shorter.
17. Capsule cylindric or tapering toward the tip, sometimes contorted, sessile or nearly so. (Subgenus *Sphaerostigma*.)
18. Flowers white (sometimes yellowish in minor and decorticans), often drying pinkish, born in terminal spikes.
19. Capsules cylindrical, terete, linear, not thickened in lower portion, scarcely if at all coiled, not noticeably attenuate at tip.
20. Petals 5-7 mm. long, suborbicular; style exceeding corolla; hypanthium 4-6 mm. long; capsule refracted or spreading. 17. Oe. refracta.
20. Petals 3 mm. long, spatulate; style shorter than corolla; hypanthium 2.5-3 mm. long; capsules divaricately spreading. . . 18. Oe. chamaenerioides.
19. Capsules not strictly cylindrical, but somewhat enlarged at the base and attenuate at tip.
21. Mature capsules usually distinctly contorted and coiled, not merely bent and curved, quite slender, not subfusiform in shape (see also Oe. decorticans var. desertorum.)
22. Flowers minute; petals 1-2 mm. long; style 1.5-3 mm. long; hypanthium 1-2 mm. long.
19. Oe. minor.

22. Flowers larger; petals 3.5-5 mm. long; style
6-12 mm. long; hypanthium 3-8 mm. long.
23. Flowers and leaves arranged in spicate tufts at
ends of naked prostrate branches or on short
central stalk; capsule 10-12 mm. long, con-
spicuously 4-angled. . . . 20. Oe. nevadensis.
23. Flowers and leaves continuous from base of
stems, not in terminal tufts; capsules 14-23
mm. long, not conspicuously 4-angled.
21. Oe. alyssoides.
21. Mature capsule merely curved or bent, not distinctly
contorted nor coiled, subfusiform in shape.
24. Leaves largely near the base, subglabrous, lance-
ovate to oblanceolate; stems glabrous or nearly
so, promptly exfoliating; capsules 15-25 mm.
long; seeds ash-colored, linear-obovoid.
22. Oe. decorticans.
24. Leaves well distributed, glandular-pubescent to
glandular-villous, ovate to oblong-ovate; stems
glandular-pubescent to glandular-villous; epi-
dermis exfoliating tardily if at all; capsule
10-15 mm. long; seeds brownish, rhomboid-pris-
matic. 23. Oe. Boothii.
18. Flowers yellow, born in axils of foliage leaves.
25. Plants with several naked, fine, often capillary stems,
each with leafy inflorescence at tip; capsule



subfusiform almost straight, 5-8 mm. long.

24. Oe. andina.

25. Plants with stems leafy from the base; capsules
terete, straight or coiled, 15-40 mm. long.

26. Flowers small; petals 2.5-3.5 mm. long; sepals 1.5-
3.5 mm. long. 25. Oe. contorta.

26. Flowers larger; petals 5-15 mm. long; sepals 3-12
mm. long. 26. Oe. dentata.

17. Capsule cylindrical or clavate, distinctly pedicelled.

(Subgenus Chylismia.)

27. Seeds oblong and with an incurving wing; plants small,
slender-stemmed, glandular-pubescent above; flowers
pinkish-white, axillary, 4-5 mm. wide.

27. Oe. pterosperma.

27. Seeds obovoid, not winged; flowers ~~not~~ axillary, but
in terminal racemes or panicles.

28. Capsules linear, elongate, usually over 2 cm. long.

29. Stems coarse, commonly branched only at base; ped-
icels short, usually 3-15 mm. long; capsules
linear, widely spreading, commonly 3-9 cm. long;
anthers hairy.

30. Stems spreading-villous, usually in lower portion;
sepals commonly pilose as well as glandular;
capsules 2-3 mm. thick. . . . 28. Oe. brevipes.

30. Stems subglabrous to finely canescent; sepals
glandular-pubescent to canescent; capsules 1-1.5
mm. thick. 29. Oe. pallidula.

1947-1948

1. The first part of the report is a general introduction to the project.

2. The second part is a description of the methods used.

3. The third part is a description of the results obtained.

4. The fourth part is a discussion of the results.

5. The fifth part is a conclusion.

6. The sixth part is a list of references.

7. The seventh part is a list of figures.

8. The eighth part is a list of tables.

9. The ninth part is a list of appendices.

10. The tenth part is a list of errata.

11. The eleventh part is a list of acknowledgments.

12. The twelfth part is a list of authors.

13. The thirteenth part is a list of titles.

14. The fourteenth part is a list of subjects.

15. The fifteenth part is a list of keywords.

16. The sixteenth part is a list of abstracts.

17. The seventeenth part is a list of summaries.

18. The eighteenth part is a list of conclusions.

19. The nineteenth part is a list of recommendations.

20. The twentieth part is a list of suggestions.

21. The twenty-first part is a list of comments.

22. The twenty-second part is a list of questions.

23. The twenty-third part is a list of answers.

24. The twenty-fourth part is a list of notes.

25. The twenty-fifth part is a list of footnotes.

29. Stems slender, commonly freely branched above; pedicels capillary, 10-25 mm. long; capsules linear, 1.5-3.5 cm. long; anthers often glabrous.

30. Oe. multijuga,

28. Capsules somewhat clavate, usually less than 2 cm. long.

31. Branches in well developed plants few to several and arising at bases of plant only, not capillary; capsules 1-2.5 cm. long; anthers scatteringly hairy; style exceeding petals.

32. Stems slender; flowers few, not congested; leaves ovate, subentire; petals less than 4 mm. long.

31. Oe. scapoidea.

32. Stems fairly coarse; flowers crowded in close terminal cluster; leaves frequently with supplementary pinnules on petioles; petals 4-7 mm. long. 32. Oe. clavaeformis.

31. Branches in well developed plants capillary and arising freely throughout the plant; anthers glabrous; style not exceeding petals.

33. Capsules 8-12 mm. long; flowers pink; anthers oblong; style pubescent at base.

33. Oe. heterochroma.

33. Capsules 3-9 mm. long; flowers yellow; anthers narrower; style glabrous. . . . 34. Oe. Parryi.

1918

1918

1918

1918

1918

1918

1918

1918

1918

1918

1918

1918

1918

1918

1918

1918

1918

1918

1. *OENOTHERA RYDBERGII* House, N. Y. State Mus. Bul. 233-234: 61.
1921.

Onagra strigosa Rydb. Mem. N.Y. Bot. Gard. 1: 278. 1900.

Oenothera strigosa Mack. & Bush, Fl. Jackson Co. Mo., 139.

1902, not of Willd. in 1825.

Biennial, grayish-strigose throughout, erect, rather simple, 3-10 dm. tall, strigose and hirsute, sometimes with reddish tinge; lowest leaves spatulate, obtuse, 3-10 cm. long, 1-2 cm. wide, with petioles 1-3 cm. long; cauline leaves lanceolate, acute, repand-denticulate, shorter petioled; inflorescence with leafy lanceolate, subsessile bracts 1-5 cm. long; hypanthium 3-4 cm. long, pubescent within, often hirsute without; sepals strigose and hirsute, 10-15 mm. long, with free tips 2 mm. long; petals yellow, 1.2-2 cm. long; stamens about as long as petals, subequal; stigma-lobes 5-6 mm. long; capsule 2.5-3.5 cm. long; seeds reddish brown, obtusely angled, irregular, 1-1.5 mm. long.

Moist places, Upper Sonoran and Transition Zones, Washington and Oregon to Minnesota. Specimen seen from Eureka Co., Nevada.

2. *OENOTHERA HOOKERI* T. & G. var. *ANGUSTIFOLIA* Gates, Mut. Factor in Evolution, 10, 30. 1915.

Biennial or possibly short-lived perennial, the stems 3-9 dm. high, reddish, somewhat muricate and hirsute, as well as strigose-canescenscent, simple or few-branched mostly from the base; leaves plane, green, the lower ones oblanceolate to spatulate, the blades 5-18 cm. long, on petioles half as long;

cauline leaves lanceolate, on shorter petioles, sinuate-dentate, gradually reduced up the stem to the leafy bracts of the inflorescence; inflorescence elongate; hypanthium 2-4 cm. long, pubescent within, reddish and with long and short hairs without; sepals red, mostly 3-5 cm. long, with long spreading hairs and short gland-tipped ones; sepal-tips 2-4 mm. long; petals 23-35 mm. long, yellow, reddish in age; capsule obtusely quadrangular, 2-5 cm. long; seeds reddish brown, sharply angled, 1.5 mm. long.

Moist places, Upper Sonoran and Transition Zones, eastern Washington and California to Colorado and New Mexico. Nevada plants seen from Washoe, Storey, Nye, Esmeralda, Mineral, Ormsby, White Pine, Churchill, Lander, Elko and Clark Counties.

3. *OENOTHERA LONGISSIMA* Rydb., Bull. Torrey Bot. Club 40: 65. 1913.

Stems 1-2 m. tall, simple or freely branched, erect, canescent, more or less hirsute, sometimes muricate; cauline leaves 8-15 cm. long, 8-15 mm. wide; bracts linear; hypanthium 8-12 cm. long; sepals 35-45 mm. long, the tips 3-6 mm. long; petals yellow, 35-45 mm. long, somewhat red in age; capsule about 4 cm. long, canescent; seeds reddish brown, 1.5-2 mm. long.

Wet places, Upper Sonoran Zone, southern Nevada and eastern California to Utah and Arizona. Nevada specimens seen from Clark Co.

4. *OENOTHERA DELTOIDES* Torr. & Frem. var. *DECUMBENS* (Wats.) Munz,

Amer. Journ. Bot. 19: 778. 1932.

Oe. albicaulis Nutt. var. decumbens S. Wats. in Parry, Amer.

Nat. 9: 270. 1875.

Oe. deltoides var. ambigua Munz, Amer. Journ. Bot. 18: 315.

1931.

Low stout annuals, simple or more frequently with central erect stems 5-25 cm. high and a few to several decumbent basal branches, these naked at base and with exfoliating epidermis; lower leaves rhombic-obovate to oblanceolate, subentire, the blades 2-7 cm. long, on equally long petioles; upper leaves coarsely sinuate-dentate; upper parts of plant with closely appressed hair; flowers solitary in upper axils; buds drooping; hypanthium slender, 2-4 cm. long, strigulose; sepals strigulose, 1.5-2 cm. long, with free tips 1-2 mm. long; petals white, aging red, 20-30 mm. long; buds 4-angled toward tip; capsules spreading, woody, with exfoliating epidermis, prismatic-cylindric to cylindric, tapering gradually from base to apex, 3-5 cm. long, 3-5 mm. thick at base; seeds narrowly obovoid, 1.5-2 mm. long, usually with purple spots.

Open deserts of Lower Sonoran Zone, southwestern Utah and southern Nevada to adjacent Arizona. Abundant in Clark County, Nevada.

var. *PIPERI* Munz, Amer. Journ. Bot. 18: 314. 1931.



Hair of upper parts of plant (hypanthia, sepals, etc.) spreading; petals usually less than 2 cm. long; capsule 1.5-3 cm. long.

Dry open places, Upper Sonoran Zone, eastern Oregon to western Nevada. Nevada plants seen from Washoe, Humboldt, Ormsby, Esmeralda and Elko Counties. (An almost glabrous form from Washoe and Ormsby Cos. has been described as forma *GLABRATA* Munz, l. c.).

5. *OENOTHERA PALLIDA* Lindl. Bot. Reg. 14: pl. 1142. 1828.

Strongly rooted perennial, with creeping rootstocks, the main stem erect, 2-5 dm. tall, branches several, spreading or ascending with white epidermis, glabrous or with few hairs in upper parts, epidermis exfoliating; leaves lanceolate to lance-linear, subentire to remotely denticulate or sinuate-dentate, usually with undulate margin, blades 2-6 cm. long, sessile or nearly so; buds acuminate, nodding; hypanthium very slender, 2-3.5 cm. long; sepals 12-18 mm. long; petals white, reddish in age, 1-3 cm. long; stamens subequal; capsule usually curved, subglabrous, 1.5-4 cm. long, subcylindric; seeds 1.5-2 mm. long, brown with dark spots or quite dark.

Sandy places and dry plains, Upper Sonoran Zone, eastern Washington to Utah and New Mexico. Nevada material seen from Lander, Elko, White Pine and Lincoln Counties.

6. *OENOTHERA CALIFORNICA* S. Wats. Bot. Calif. 1: 223. 1876.

Perennial from underground rootstocks, rather coarse-stemmed, usually branched, 1-4 dm. tall, frequently decumbent or ascending, ashy with short appressed hairs throughout and with some longer spreading ones especially in upper parts, epidermis exfoliating; leaves variable, blades of lower ones oblanceolate to spatulate in outline, of cauline ones oblong to lanceolate, from subentire to deeply and regularly sinuate-dentate, 1-6 cm. long, sessile or nearly so; flowers several, buds nodding; hypanthium slender, 2-4 cm. long, strigulose and villous; sepals lance-linear, 1.5-2 cm. long, with free tips nearly or quite wanting; petals 2-3 cm. long; stamens subequal; capsule terete, usually divaricate and somewhat curved upward, 2-5 cm. long; seeds 1.5 mm. long, brown with dark spots.

Dry plains of Upper Sonoran and Lower Transition Zones, eastern and southern California, adjacent Nevada, Lower California. Nevada plants seen from Esmeralda, Lander, Nye and Clark Counties.

7. *OENOTHERA CAESPITOSA* Nutt. var. *MARGINATA* (Nutt.) Munz, Amer.

Journ. Bot. 18: 733. 1931.

Oe. marginata Nutt. ex H. & A. Bot. Beechey, 342. 1838.

Caespitose perennial, acaulescent, or more usually caulescent, villous-hirsute throughout; leaves oblanceolate, usually

sinuate-pinnatifid, the blades 3-10 cm. long, on winged petioles nearly as long; flowers fragrant, vespertine, white, aging red; hypanthium 5-8 cm. long, often tinged red; sepals 2.5-3.5 cm. long, hairy, scarcely free-tipped; petals broadly obovate, 2.5-4 cm. long; stamens subequal, glabrous; stigma-lobes 5-8 mm. long; capsule pedicelled, linear-cylindric, straight, 3-4 cm. long, scarcely ridged, with low tubercles; seeds dark brown, 3 mm. long, obovoid, minutely cellular-roughened, conspicuously furrowed along the raphe.

Dry slopes, Upper Sonoran and Transition Zones, eastern Washington and California to Montana and New Mexico. Practically throughout Nevada at elevations between 3000 and 9000 feet.

var. MONTANA (Nutt.) Durand, Bot. Basin of Great Salt Lake of Utah,

164. 1859.

Oe. montana Nutt. ex Torr. & Gray, Fl. No. Amer. 1: 500.

1840.

Plant acaulescent; leaves usually shorter and wider than in preceding, canescent-pubescent on margins and often on veins of lower surface, otherwise quite glabrous; capsule sessile, ovoid, not tubercled, ca. 2 cm. long.

Dry slopes, Upper Sonoran Zone, eastern Oregon and California to Colorado and Nebraska. Nevada material seen from Humboldt and Elko Counties.

var. PURPUREA (Wats.) Munz, Amer. Jour. Bot. 18: 730. 1931.

Oe. marginata var. purpurea S. Wats., Bot. King Exped. 108.
1871.

Pachylophus canescens Piper, Contr. U. S. Nat. Herb. 11:
409. 1906.

Acaulescent, densely canescent throughout with a fine appressed pubescence; capsule 1-2 cm. long, with low rounded tubercles on angles.

Dry slopes, Upper Sonoran and Transition Zones, eastern Washington and California to Montana and Wyoming. Nevada material seen from Lander, Elko and White Pine Counties.

var. LONGIFLORA (Heller) Munz, Amer. Jour. Bot. 18: 734. 1931.

Anogra longiflora Heller, Muhlenbergia 2: 224. 1906.

Habit of var. marginata, but subglabrous except for a few hairs along the margins of the leaves, about the ovaries and sepals which latter may also be finely glandular-puberulent.

Inyo County, California and Esmeralda Co., Nevada.

var. CRINITA (Rydb.) Munz, Amer. Jour. Bot. 18: 731. 1931.

Pachylophus crinitus Rydb. Fl. Rocky Mts. & Adj. Plains,
598 & 1064. 1917.

Caespitose from a woody caudex; stems 3-8 cm. high; leaves densely hirsute or even pilose, crowded, the blades lanceolate, obtuse, sinuate, 1-1.5 cm. long, 3-5 mm. wide; hypanthium 2-4

cm. long, hirsute; capsules 1-1.4 cm. long, 3-5 mm. thick.

At high altitudes, southern Utah and Nevada, from which latter state material has been seen from White Pine, Nye and Clark Counties.

8. *OENOTHERA CAVERNAE* Munz. Leaflets West. Bot. 3: 50. 1941.

Acaulescent, caespitose, winter annual; leaves in a rosette, regularly lyrate-pinnatifid into simple lobes or these somewhat crenate-dentate, leaf-blades oblanceolate in outline, glandular-puberulent, pilose especially on margins and veins, 2.5-10 cm. long, 0.6-2.0 cm. wide; petioles one-third as long; hypanthium slender, whitish, pilose and glandular-puberulent, 2.5-4 cm. long; sepals 5-9 mm. long; petals white, scarcely red in age, 8-12 mm. long; style and stamens slightly more than half the petals; capsule sessile, woody, lance-ovoid, 1.5-3 cm. long, 0.6-1.0 cm. thick, somewhat pilose, with 8 longitudinally tuberculate ridges; seeds brown, 2 mm. long, with furrowed raphe.

On limestone, from 3 localities in Clark Co., Nevada: Gypsum Cave, foothills of Spring Mts. above Arden, and Sheep Mt. above Jean.

9. *OENOTHERA PRIMIVERIS* Gray, Pl. Wright. 2: 58. 1853.

Lavauxia lobata A. Nels., Bot. Gaz. 47: 429. 1909.

Annual or winter annual, with long tap-root, caespitose, acaulescent or nearly so, villous or pilose throughout;

leaf-blades oblanceolate in outline, 1-12 cm. long, usually deeply and regularly pinnatifid into lanceolate or ovate lobes which are in turn lobed or toothed; petioles shorter than blades; flowers vespertine, yellow, aging orange-red; hypanthium 2-6 cm. long; sepals lance-linear, 15-28 mm. long, without free tips; petals cuneate-obovate, usually 2-3 cm. long, with terminal sinus; stamens subequal; stigma lobes 6-8 mm. long; capsule pilose, quadrangular, with heavy rib down middle of each face, reticulate, not winged nor tuberculate, gradually tapering to attenuate apex, 2-3.5 cm. long, 6-8 mm. thick at base; seeds brown, somewhat roughened-tuberculate, 2.5-3 mm. long, with narrow raphal groove.

Dry plains, Lower Sonoran Zone, deserts from California to Utah, and Texas. Seen from Nye and Clark Counties, Nevada.

10. *OENOTHERA XYLOCARPA* Coville, Contr. U. S. Nat. Herb. 4: 105.
pl. 8. 1893.

Acaulescent perennial, with thick caudex and crown of leaves; leaf-blades pinnately parted, often spotted red, 2-7 cm. long, broadly oblanceolate to obovate in outline, with dense soft sometimes grayish pubescence, the terminal lobe much the largest; petioles about as long as blades; hypanthium slender, villous, 2.5-4.5 cm. long; sepals 2-3 cm. long; petals bright yellow, aging salmon-red, 2.5-3 cm. long; stamens subequal; stigma-lobes 4-5 mm. long; capsules somewhat woody, 3.5-6 cm. long, the body proper 7-8 mm. thick at base and winged,

tapering gradually into a long slender wingless upper portion, capsule 4-faced, with median nerve on each face; seeds brown, 2-2.5 mm. long, narrowly obovoid, angled, roughened and minutely tuberculate, with a broad flat raphe.

Dry benches among pines, Transition Zone, east central California and Washoe County, Nevada.

11. *CENOTHERA FLAVA* (A. Nels.) Garrett, Spring Flora of Wasatch region, 4th edition, 106. 1927.

Lavauxia flava A. Nels. Bull. Torrey Bot. Club 31: 243.
1904.

Perennial, with thick taproot, acaulescent or nearly so, subglabrous to finely glandular or pubescent especially about the flowers; leaf-blades oblong-linear to oblanceolate in outline, 3-20 cm. long, 1-2 cm. wide, deeply and irregularly runcinate-pinnatifid, with a broadly winged rachis; petiole shorter; hypanthium slender, 2-12 cm. long; sepals 10-16 mm. long; petals yellow, 1-2 cm. long; stamens subequal; stigma-lobes 3-4 mm. long; capsule indurate, ovoid, 1-2 cm. long, 4-winged, each wing reticulate-veined, 2-5 mm. wide especially above; seeds numerous, dark brown, 2 mm. long, slightly concave with carinate ridge on ventral side and wing-like margin around obtuse summit.

About desiccating depressions, high Upper Sonoran and Transition Zones, northern California and eastern Washington

to Rocky Mts. Nevada material seen from Washoe, Douglas, Ormsby, Lyon, Elko and White Pine Counties.

12. *OENOTHERA BRACHYCARPA* Gray var. *WRIGHTII* (Gray) Lévl. Mon.

Onoth. 40. 1902.

Oe. Wrightii Gray, *Pl. Wright.* 2: 57. 1853.


Lavauxia Howardi (Jones) Nelson, *Bot. Gaz.* 34: 368. 1902.

Subcaulescent perennial, with heavy caudex, closely cinereous-strigulose or even hoary throughout; leaves tufted, thick in texture, lanceolate to lance-ovate in outline, greenish or hoary, entire to sinuate-pinnatifid or even with a few lateral lobes, the blades 4-15 cm. long, 1-3 cm. wide; petioles about as long; hypanthium 5-15 cm. long; sepals 3-5 cm. long, with free tips 1-4 mm. long; petals yellow, drying reddish, 3-5 cm. long; stigma-lobes 5-10 mm. long; capsules 2.5-3 cm. long, 6-8 mm. thick, ovoid to cylindric-ovoid, winged from only near the summit to almost the entire length, the wings 1.5-4.5 mm. wide; seeds dark brown, more or less corky tubercled, the summit with denticulate corky wing.

Dry places, apparently often on limestone, Sonoran and Transition Zones, Idaho to Colorado, Arizona and Mexico. Nevada material seen from Lincoln and Clark Counties.

13. *OENOTHERA LAVANDULAEFOLIA* T. & G. var. *GLANDULOSA* Munz, *Amer.*

Journ. Bot. 16: 705. 1929.



Low caespitose, suffrutescent perennials from woody caudex, 5-20 cm. tall; stems and foliage grayish-strigose throughout; leaves linear to narrowly oblanceolate, 5-15 (25) mm. long, 1-3 mm. wide, sessile or nearly so, crowded on stems, not reduced up the stem; flowers few, solitary in upper axils; hypanthium 2.5-5 cm. long, glandular-pubescent; sepals 8-12 mm. long, glandular-pubescent; petals yellow, reddish in age, rhomboidal, 13-22 mm. long; stamens subequal, two-thirds the petals; stigma discoid, squarish, 2-4 mm. wide; capsule cylindrical, often somewhat fusiform or clavate, 8-20 mm. long, 2-3 mm. thick; seeds brownish, obovoid, angled, 1 mm. or more long.

Dry slopes and ridges, apparently usually in limestone areas, Transition Zone, Nevada to Colorado and Texas. Nevada plants seen from White Pine, Lincoln and Clark Counties.

14. *OENOTHERA PALMERI* S. Wats. Proc. Amer. Acad. 12: 251. 1877.

Taraxia Palmeri Small, Bull. Torrey Bot. Club 23: 184. 1896.

Dwarf caespitose annual, with slender taproot, finely strigose throughout, forming small caulescent tufts 2-6 cm. tall, or with several short horizontal branches 2-4 cm. long; stems pubescent; leaves linear-lanceolate to -oblanceolate, subentire, 2-6 cm. long; sterile portion of upper part of ovary filiform, 8-18 mm. long; hypanthium proper (above sterile portion of ovary) 1-2 mm. long; sepals 2-3 mm. long; petals yellow, 3-5 mm. long; capsules crowded, ovoid, 5-7 mm. long,

coriaceous and tough, 4-angled below, each angle growing into an obliquely truncate wing along the upper edge of which is the line of dehiscence; seeds smooth, brownish, 1.5 mm. long.

Open places, deserts of Sonoran Zones, eastern Oregon to California and Arizona. Nevada material seen from Ormsby Co.

15. *OENOTHERA HETERANTHA* Nutt. ex T. & G. Fl. No. Amer. 1: 507.

1840.

Oe. heterantha var. taraxacifolia S. Wats., Proc. Amer. Acad.

8: 589. 1873.

Oe. subacaulis Garrett, Spring Fl. Wasatch Reg. 64. 1911.

Acaulescent perennial, essentially glabrous, except for some pubescence on leaf-margins; leaf-blades lanceolate to ovate-lanceolate, entire to sinuate, 3-15 cm. long, 1-5 cm. wide, narrowed into equally long petioles; ovary with an upper sterile slender portion 3-10 cm. long; sepals 5-8 mm. long; petals yellow, 8-10 mm. long; stigma discoid, 1.5-2 mm. wide; capsule oblong-ovoid, relatively smooth, coriaceous, persistent, somewhat 4-angled, 12-15 mm. long; seeds oblong, straw-colored, 3 mm. long.

Moist grassy places, Upper Sonoran and Transition Zones, eastern Washington to California and Rocky Mts. Nevada material seen from Washoe, Humboldt, Douglas, Lander, Elko and White Pine Cos.

16. *OENOTHERA TANACETIFOLIA* Torr. & Gray, Pac. R.R.Rep. 2: 121.

pl. 4. 1854.

Taraxia tanacetifolia Piper, Contr. U.S. Nat. Herb. 11: 405.

1906.

Perennial, with thick root and simple or branched crown, subglabrous to densely and finely pubescent; leaves lanceolate in outline, deeply sinuate-pinnatifid, the blades 3-10 cm. long, the numerous segments unequal, acute, entire or toothed, the petioles about as long as the blades; sterile portion of the ovary pubescent, slender, 2-10 cm. long; hypanthium proper 3 mm. long; sepals 7-9 mm. long; petals yellow, aging red, 10-15 mm. long; stamens unequal; stigma globular; capsule rather rarely developed, pubescent, narrowly ovoid, quadrangular, torulose, 17-20 mm. long, 5-6 mm. thick, relatively straight; seeds numerous, brown, carunculate, 2 mm. long.

Damp grassy places, Upper Sonoran and Transition Zones, eastern Washington to Mono Co., California, Idaho and Nevada. Nevada material seen from Washoe, Ormsby, Douglas and Elko Counties.

17. *OENOTHERA REFRACTA* S. Wats. Proc. Amer. Acad. 17: 373. 1882.

Sphaerostigma refractum Small, Bull. Torrey Bot. Club

23: 192. 1896.

Annual, 5-40 cm. tall, erect, with few to several divaricately spreading branches, usually glandular-puberulent and

somewhat strigulose throughout, the stems slender, commonly reddish, with exfoliating epidermis; leaves largest near base of plant, but well distributed to lowest flowers, oblanceolate to lanceolate to oblong-linear, entire to denticulate, 2-5 cm. long, short-petioled to sessile; inflorescence racemose to paniculate, 5-15 cm. long; hypanthium 5-6 mm. long; sepals 5-6 mm. long; petals white, suborbicular, 4-7 mm. long; stigma globose; capsule linear, commonly refracted or spreading, 3-5 cm. long, straight to somewhat coiled; seeds pale, linear, 1 mm. long.

Open places, deserts of Lower Sonoran Zone, California to Utah and Arizona. Nevada material seen from Nye and Clark Counties.

18. *OENOTHERA CHAMENERIOIDES* Gray, Pl. Wright. 2: 58. 1853.

Sphaerostigma chamaenerioides Small, Bull. Torrey Bot.

Club 23: 189. 1896.

Erect annual, 1-5 dm. tall, usually branching near the base, stems slender, often reddish, lower portions glandular-puberulent, upper strigulose and glandular-puberulent; leaf-blades thin, 4-8 cm. long, ovate-lanceolate to lanceolate, entire, with petioles 1-3 cm. long; inflorescence a corymbose raceme, elongating in fruit to nearly 2 dm; hypanthium 2.5-3 mm. long; sepals 2.5 mm. long; petals white, often reddish in age, 3 mm. long; capsule terete, linear, divaricately spreading,

scarcely if at all beaked, 25-50 mm. long; seeds linear, 1 mm. long.

Open places, deserts of Lower Sonoran Zone, California to Utah, Texas, and Mexico. For Nevada, seen from Nye, Lincoln and Clark Counties, and possibly Washoe and Ormsby Counties.

19. *OENOTHERA MINOR* (A. Nels.) Munz, Bot. Gaz. 85: 238. 1928.

Sphaerostigma minus A. Nelson, Bull. Torrey Bot. Club
26: 130. 1899.

S. tortum A. Nels. Bot. Gaz. 40: 60. 1905.

Annual, more or less canescent-strigulose throughout, scarcely if at all glandular; stems simple and erect, or usually with several subequal ascending stems, 5-20 cm. high; basal leaves largest, blades spatulate to oblanceolate to elliptic-ovate, subentire, 5-25 mm. long, 2-15 mm. wide, narrowed gradually into petioles 5-20 mm. long; upper leaves narrower and smaller; flowers borne singly in axils of almost all leaves, the upper in a spicate inflorescence; hypanthium 2 mm. long; sepals and petals 2 mm. long; capsules 13-25 mm. long, more or less contorted, often beaked; seeds narrowly obovoid, 1 mm. long.

Dry slopes, Upper Sonoran Zone, Idaho to Wyoming and Nevada. Nevada material seen from Pershing, Lander, Eureka, Elko and White Pine Counties.

20. *OENOTHERA NEVADENSIS* Kellogg, Proc. Calif. Acad. 2: 224. fig.

70. 1863.

Sphaerostigma nevadense Heller, *Muhlenbergia* 6: 51. 1910.

S. tortuosum A. Nels. *Proc. Biol. Soc. Wash.* 17: 95. 1904.

Annual, subglabrous, forming a simple erect tuft, 2-5 cm. tall, or with several naked prostrate branches 3-10 cm. long with terminal tufts of leaves and flowers (these tufts may elongate in fruit); leaves narrowly oblanceolate, 1-3.5 cm. long, on petioles of about same length; flowers white; petals 3-5 mm. long; capsules 10-12 mm. long, 1.5-2 mm. thick, quadrangular with ridge along middle of each face, swollen at base, narrowed toward slender beak, coiled and twisted, usually crowded; seeds pale gray, 1 mm. long.

Depressions, Upper Sonoran Zone, Washoe, Ormsby, and Storey Counties, Nevada.

21. *OENOTHERA ALYSSOIDES* H. & A. var. *VILLOSA* S. Wats. *Proc. Amer. Acad.* 8: 591. 1873.

Sphaerostigma macrophyllum Rydb. *Bull. Torrey Bot. Club* 40: 66. 1913.

S. utahense Small, *Bull. Torrey Bot. Club* 23: 191. 1896.

Annual, usually branching from base, central stem erect, others ascending and curved at tip, grayish-canescens or grayish-villous, 5-35 cm. tall; leaves grayish, oblanceolate to ovate-lanceolate, 15-40 mm. long, entire or nearly so, the lowermost petioled, upper sessile; hypanthium 4-8 mm. long; sepals 4-5 mm. long; petals white, aging reddish, sometimes drying yellowish, 4-5 mm. long; capsules 15-25 mm. long,

thickened at base, gradually attenuate toward beaklike tip, much contorted or only curved; seeds pale, linear-obovoid.

Dry plains and washes, Upper Sonoran Zone, Utah to eastern California and Arizona. In Nevada growing almost throughout the state at elevations of 4000 to 7000 feet.

22. *OENOTHERA DECORTICANS* (H. & A.) Greene var. *CONDENSATA* Munz, Bot. Gaz. 85: 247. 1928.

Annual, erect, simple or branching below, the branches ascending or spreading, nearly glabrous except in the finely pubescent and sometimes glandular inflorescence, stems thick, 1-2 dm. tall, the pure white epidermis exfoliating; leaves in the lower part, subentire, 2-6 cm. long, with equally long petioles, lanceolate to oblanceolate, uppermost reduced; inflorescence a compact spike; hypanthium 4-6 mm. long; sepals 4-5 mm. long; petals white, aging reddish, suborbicular, 4-5 mm. long; capsules woody, much thickened, about 3 mm. at base, 15-25 mm. long, beaked, simply curved, so that the beak points away from the stem; seeds ash-colored, linear-obovoid, 1 mm. long.

The persistent dead stalks and cone-like fruiting spikes of woody capsules are a prominent feature of the open deserts of California and adjacent regions, ranging to so. Utah and Arizona. Nevada material seen from Nye and Clark Counties.

var. *DESERTORUM* Munz, Bot. Gaz. 85: 246. 1928.

More slender-stemmed; petals longer than wide; capsules 1-1.5 mm. thick.

Lower Sonoran Zone of California deserts; collected at Rhyolite, Nye Co., Nevada.

23. *CENOTHERA BOOTHII* Dougl. ex Hook. Fl. Bcr. Amer. 1: 213. 1834.

Sphaerostigma Boothii Walp. Rep. 2: 77. 1843.

S. senex A. Nels. Proc. Biol. Soc. Wash. 18: 173. 1905.

Annual, glandular-pubescent to glandular-villous throughout, erect, 1-4 dm. tall, usually with central stem more prominent than the spreading branches; leaves ovate to oblong-ovate, well distributed, subentire, 2-5 cm. long, with petioles 1-3 cm. long; inflorescence racemose-spicate, often quite congested, elongating in fruit; hypanthium 4-8 mm. long; sepals 3-7 mm. long; petals white, pinkish in age, obovate, clawed, 4-9 mm. long; stamens subequal; capsule 10-15 mm. long, usually ascending in lower half and with terminal portion spreading but not contorted, thickest near base, 1.5-2 mm. thick; seeds brown, rhomboid-prismatic, 1 mm. long.

Dry plains and slopes, especially in gravelly and disturbed places, Upper Sonoran Zone, eastern Washington and California to Utah and Arizona. Nevada material seen from Washoe, Ormsby, Churchill, Lander, Mineral, Esmeralda, and Nye Counties.

24. *CENOTHERA ANDINA* Nutt. ex Torr. & Gray, Fl. No. Amer. 1: 512.

1840.

Sphaerostigma andinum Walp. Rep. 2: 79. 1843.


Low, erect, very slender-stemmed annuals, with spreading branches from near the base or above, finely canescent throughout, 2-15 cm. tall and about as broad, lower stem and branches in all except the smallest plants rather free of leaves; leaves alternate, linear to narrowly oblanceolate, entire, with short indistinct petioles, blades 1-3 cm. long; flowers axillary in a rather crowded corymb which becomes racemose in fruit; hypanthium 1 mm. long; sepals 1.5 mm. long; petals yellow, 1.5 mm. long; capsule 5-6 mm. long, fusiform, somewhat quadrangular; seeds fusiform, smooth, brown, 0.7 mm. long.

Dried depressions, plains, Upper Sonoran Zone, eastern Washington to northeastern California, Assiniboia and Utah. Nevada material seen from Humboldt, Eureka and Elko Counties.

25. OENOTHERA CONTORTA Dougl. ex Hook. Fl. Bor. Amer. 1: 214. 1834.

Sphaerostigma contortum Walp. Rep. 2: 78. 1843.

Annual, slender-stemmed, 5-10, or more cm. tall, subglabrous to finely pubescent, usually with several to few suberect branches from near the base; leaves well distributed, linear to lance-linear, not more than 2 mm. wide, 5-25 mm. long, subsessile, upper ones reduced to leafy bracts and subtending the flowers; hypanthium 1-2 mm. long; sepals 1.5-2.5 mm. long; petals bright yellow, aging red, narrowly obovate to obcordate, 2.5-3 mm. long; capsules linear, cylindrical, often torulose, sessile, curved or straight, 25-35 mm.



long, ending in a definite beak; seeds brown, obovoid, less than 1 mm. long.

Dry loose slopes, recently disturbed places, sandy areas, etc. of Upper Sonoran Zone, eastern British Columbia to eastern California, and Nevada. Nevada plants seen from Washoe, Ormsby and Nye Counties.

var. FLEXUOSA (A. Nels.) Munz, Bot. Gaz. 85: 253. 1928.

Sphaerostigma flexuosum A. Nels. ex Rydb. Fl. Rocky Mts. 601. 1917.

Stems and leaves as in the species; capsules more slender, distinctly pedicelled, not beaked, 17-25 mm. long, frequently curved into a half circle.

Ranging with the species but extending further south and east. Nevada material seen from Washoe, Humboldt, Mineral, Esmeralda and Elko Counties.

var. PUBENS (S. Wats.) Coville, Contr. U. S. Nat. Herb. 4: 104. 1893.

Oe. strigulosa var. pubens S. Wats. Proc. Amer. Acad. 8: 594. 1873.

Sphaerostigma orthocarpum Nels. & Kennedy, Proc. Biol. Soc. Wash. 19: 155. 1906.

Coarse-stemmed plants with abundant spreading pubescence; leaves commonly over 2 mm. wide; capsules 1 mm. or more in diameter, 25-36 mm. long, sessile or subsessile, not beaked.

Eastern middle California and western Nevada. Material

seen from Washoe, Storey, Ormsby, Douglas, Lander, Mineral, and Nye Counties, Nevada.

26. *OENOTHERA DENTATA* Cav. var. *JOHNSTONII* Munz, Bot. Gaz. 85:

259. 1928.

Annual, usually branched from base, stems subdecumbent to ascending, subglabrous to glandular-pubescent, slender, 5-25 cm. tall, with light colored epidermis inclined to exfoliate; leaves mostly lance-linear, subsessile, 1-3 cm. long; flowers few, solitary in upper axils; inflorescence glandular-pubescent; sepals 5-12 mm. long; petals yellow, 10-16 mm. long; capsules linear, terete, somewhat torulose, straight to somewhat contorted, 3-4 cm. long, 1 mm. thick, not conspicuously beaked; seeds 1 mm. long.

Lower Sonoran Zone, California deserts. Known in Nevada from a single collection at Nipton, Clark County. [This locality is really in California, but is given on M. E. Jones labels as Nevada. Where the plants were actually collected is not known.]

27. *OENOTHERA PTEROSPERMA* S. Wats. Bot. King, 112. pl. 14. 1871.

Chylismia pterosperma Small, Bull. Torrey Bot. Club 23:

193. 1896.

Annual, low, 5-12 cm. tall, erect, simple or with few open branches; stems slender, finely pilose below, finely glandular above; leaves oblong- to ovate-lanceolate, entire, sessile or nearly so, 5-20 mm. long; flowers axillary, pinkish white;

1870

1871

1872

1873

1874

1875

1876

1877

1878

1879

1880

1881

1882

1883

1884

1885

1886

1887

1888

1889

1890

1891

1892

1893

1894

1895

1896

1897

1898

1899

1900

1901

1902

1903

1904

1905

1906

1907

1908

1909

1910

1911

1912

1913

1914

1915

1916

1917

1918

1919

1920

1921

1922

1923

1924

1925

1926

1927

1928

1929

1930

1931

1932

1933

1934

1935

1936

1937

1938

1939

1940

1941

1942

1943

1944

1945

1946

1947

1948

1949

1950

1951

1952

1953

1954

1955

1956

1957

1958

1959

1960

1961

1962

1963

1964

1965

1966

1967

1968

1969

1970

1971

1972

1973

1974

1975

1976

1977

1978

1979

1980

1981

1982

1983

1984

1985

1986

1987

1988

1989

1990

1991

1992

1993

1994

1995

1996

1997

1998

1999

2000

2001

2002

2003

2004

2005

2006

2007

2008

2009

2010

2011

2012

2013

2014

2015

2016

2017

2018

2019

2020

2021

2022

2023

2024

2025

2026

2027

2028

2029

2030

2031

2032

2033

2034

2035

2036

2037

2038

2039

2040

2041

2042

2043

2044

2045

2046

2047

2048

2049

2050

2051

2052

2053

2054

2055

2056

2057

2058

2059

2060

2061

2062

2063

2064

2065

2066

2067

2068

2069

2070

2071

2072

2073

2074

2075

2076

2077

2078

2079

2080

2081

2082

2083

2084

2085

2086

2087

2088

2089

2090

2091

2092

2093

2094

2095

2096

2097

2098

2099

2100

2101

2102

2103

2104

2105

2106

2107

2108

2109

2110

2111

2112

2113

2114

2115

2116

2117

2118

2119

2120

2121

2122

2123

2124

2125

2126

2127

2128

2129

2130

2131

2132

2133

2134

2135

2136

2137

2138

2139

2140

2141

2142

2143

2144

2145

2146

2147

2148

2149

2150

2151

2152

2153

2154

2155

2156

2157

2158

2159

2160

2161

2162

2163

2164

2165

2166

2167

2168

2169

2170

2171

2172

2173

2174

2175

2176

2177

2178

2179

2180

2181

2182

2183

2184

2185

2186

2187

2188

2189

2190

2191

2192

2193

2194

2195

2196

2197

2198

2199

2200

pedicels 5-8 mm. long, capillary; hypanthium 1.2 mm. long; sepals 1.5-2.5 mm. long; petals obcordate, 1.5-2.5 mm. long; stamens unequal; capsules cylindric-clavate, slightly curved, erect, 10-16 mm. long, attenuate at base; seeds oblong, 1.5 mm. long, brownish, flattened, bordered with a revolute, wing-like margin which is covered with cellular processes.


Uncommon, in dry places, Upper Sonoran Zone, eastern Oregon to California and Utah. Nevada plants seen from Washoe, Pershing, Esmeralda, Elko and Clark Counties.

28. *OENOTHERA BREVIFES* Gray, Pac. R. R. Rep. 4: 87. 1857.

Chylismia brevipes Small, Bull. Torrey Bot. Club 23: 194, 1896.

Annual, frequently rather coarse, usually 1- to few-stemmed from base, not much branched above, spreading-villous, 1-4 dm. tall, erect with nodding stem-tips; leaves largely in basal rosette, petioled, subglabrous to villous, ovate to oblong-cordate, subentire to pinnatifid, 3-8 cm. long, on petioles equally long, uppermost leaves few and much reduced; inflorescence racemose; pedicels 3-15 mm. long; hypanthium 3-7 mm. long with pubescent papillate structure on each rib within; sepals 6-10 mm. long; petals bright yellow, diurnal, 7-15 mm. long; capsule linear, spreading-divaricate, 5-9 cm. long, 2-3 mm. thick; seeds straw-colored, obovoid, 1-1.5 mm. long.

Common in washes, open deserts of Lower Sonoran Zone from California to Nevada and Arizona. Nevada plants seen from Clark Co.



29. OENOTHERA PALLIDULA Munz, Leaflets West. Bot. 2: 88. 1938.

Oe. brevipes var. pallidula Munz, Amer. Jour. Bot. 15:

229. 1928.

With habit of the preceding species, but stems ashy-strigose, not spreading-villous; flowers paler yellow; hypanthium lacking a swollen structure within on each rib at upper end of the internal pubescence; sepals strigulose to subglabrous, not pilose; petals 8-12 mm. long; capsules 2-5 cm. long, 1-2 mm. thick.

Lower Sonoran Zone from Death Valley region of California to southwestern Utah, Arizona. Nevada plants seen from Nye and Clark Counties.

30. OENOTHERA MULTIJUGA S. Wats. Amer. Nat. 7: 300. 1873.

Chylismia multijuga Small, Bull. Torrey Bot. Club 23:

193. 1896.

C. venosa Nels. & Kennedy, Muhlenbergia 3: 140. 1908.

C. hirta A. Nels. Bot. Gaz. 47: 428. 1909.

Annual to biennial, subglabrous to pubescent to villous, slender, one-stemmed and erect, or branching freely from the base, freely branched above, 2-8 dm. tall; leaves mostly in a basal rosette, 10-25 cm. long, usually much pinnate, with 6-20 pairs of major pinnae and smaller ones often in between, terminal pinna often conspicuously larger; petioles 2-6 cm. long; inflorescence of several naked racemes in a loose open panicle 1-6 dm. long; pedicels capillary, 1-2.5 cm. long;

hypanthium funnellform; sepals 6-7 mm. long, with or without free tips; petals yellow, 7-9 mm. long; pedicels capillary, 7-15 mm. long; capsules slender, linear, 15-35 mm. long, 1-1.5 mm. thick; seeds light brown, obovoid, 1 mm. long.

Nevada to Utah and Arizona, growing in washes and disturbed places of the Lower Sonoran Zone. Nevada plants seen from Clark County.

var. PARVIFLORA (S. Wats.) Munz, Amer. Journ. Bot. 15:

231. 1928.

Oe. brevipes var. parviflora S. Wats. ex Parry, Amer. Nat.

9: 271. 1875.

Leaflets often fewer; flowers smaller; sepals 3-4 mm. long; petals 3-5 mm. long.

Lower Sonoran deserts, eastern California to Utah. Nevada plants seen from Esmeralda, Nye, Clark, Lincoln and Elko Counties.

31. OENOTHERA SCAPOIDEA Nutt. ex Torr. & Gray, Fl. No. Amer. 1:

506. 1840.

Chylismia scapoidea Small, Bull. Torrey Bot. Club 23:

193. 1896.

Annual, simple or branching from base, erect or spreading, glabrous, the stems slender, simple or nearly so, 1-4.5 dm. high; leaves mostly basal, mostly simple, ovate to oblong-ovate, subentire to dentate, the blades 1-4 cm. long, petioles 2-7 cm. long; inflorescence mostly racemose; pedicels capillary,



5-15 mm. long; hypanthium 1.5-3 mm. long; sepals 2-3 mm. long; petals yellowish, often red-dotted at base, 2-4 mm. long; capsules erect, clavate, slightly curved, 1-2.5 cm. long, 2-2.5 mm. thick; seeds brownish, obovoid, 1.5-2 mm. long.

Dry slopes and exposed places, Upper Sonoran Zone, Nevada to Wyoming and Utah. Nevada material seen from Humboldt and Elko Counties.

32. *OENOTHERA CLAVAEFORMIS* Torr. & Frém. in Frém. Rep. 314. 1845.

Chylismia clavaeformis Heller, *Muhlenbergia* 2: 105. 1906.

Annual, simple or with few unbranched stems from the base, 1-4 dm. tall, subglabrous to finely pubescent below, subglabrous to glandular in the inflorescence; leaves mostly in a basal rosette, simple and irregularly dentate, with ovate blades 2-5 cm. long and petioles of same length, rarely pinnatifid; upper leaves much reduced; inflorescence racemose, somewhat peduncled, the flowers quite crowded at anthesis; pedicels 8-25 mm. long; sepals 5 mm. long; petals white, often drying reddish, 4-6 mm. long; capsule clavate, commonly 2 mm. thick, 12-20 mm. long, generally curved and ascending; seeds light brown, obovoid, 1.2 mm. long.

Dry plains, deserts of Upper and Lower Sonoran Zones, California to Nevada. Seen from following Nevada Counties: Washoe, Humboldt, Storey, Pershing, Churchill, Lander, Eureka, Elko, Lyon, Mineral, Esmeralda and Nye.



var. AURANTIACA (S. Wats.) Munz, Amer. Journ. Bot. 15:

237. 1928.

Oe. scapoidea var. aurantiaca S. Wats. Proc. Amer. Acad.

8: 595. 1873.

Hypanthium and sepals strigulose; flowers sometimes more yellowish, even with orange in hypanthium.

Lower Sonoran Zone, deserts of California to Utah and Arizona. Nevada material seen from Nye and Clark Counties.

var. PURPURASCENS (S. Wats.) Munz, Leaflets West. Bot. 3:

53. 1941.

Oe. scapoidea var. purpurascens S. Wats. Proc. Amer. Acad.

8: 595. 1873.

Oe. claviformis var. cruciformis (Kell.) Munz, Amer. Journ.

Bot. 15: 235. 1928.

Stems puberulent; leaves scarcely or not pinnatifid; flowers yellow, sometimes with reddish spots.

Loose dry slopes and disturbed areas, Upper Sonoran Zone, eastern Oregon to eastern California and adjacent Nevada, for which it has been seen from the following Counties: Washoe, Storey, Ormsby, Pershing and Churchill.

33. OENOTHERA HETEROCHROMA S. Wats. Proc. Amer. Acad. 17: 373.

1882.

Chylismia heterochroma Small, Bull. Torrey Bot. Club 23:

193. 1896.

Annual, simple or branched at base, branched above, gland-



ular-pubescent throughout, 2-5 dm. tall; leaves in lower portion only, but not actually in basal rosette, ovate, irregularly serrate, villous, 2-6 cm. long, on petioles almost as long; upper leaves reduced; pedicels capillary, 2-5 mm. long; hypanthium 2.5 mm. long; sepals 2.5 mm. long; petals purplish, 3-5 mm. long; stamens unequal; capsules 8-13 mm. long, 2 mm. thick, clavate; seeds brown, obovoid, 1 mm. long.

Uncommon, dry places, Upper Sonoran Zone, eastern California to Nevada. Nevada localities in Lander, Mineral, Esmeralda and Lincoln Counties.

var. *MEGALANTHA* Munz, Leaflets West. Bot. 3: 52. 1941.

Hypanthium 7 mm. long; petals 1 cm. long.

Known from a single collection: Train 2358 from 50 miles southeast of Beatty, Skull Mts., Nye Co., Nevada.

34. *OENOTHERA PARRYI* S. Wats. Amer. Nat. 9: 19 & 270. 1875.

Chylismia Parryi Small, Bull. Torrey Bot. Club 23: 193.

1896.

Oenothera tenuissima M. E. Jones, Proc. Calif. Acad., ser.

2, 5: 682. 1895.

Annual, 1-3 dm. high, erect, mostly branched, lower portion villous with spreading hairs; lower leaves crowded or well distributed, lanceolate to oblong-ovate, villous beneath, subglabrous above, 1-2.5 cm. long, subentire to subsinuate-toothed, the petioles 1-3.5 cm. long; uppermost leaves reduced

to ovate bracts; inflorescence paniculate, with numerous fine, glandular-pubescent branches; pedicels capillary, recurved, 5-16 mm. long; hypanthium 1-2 mm. long; sepals 2-4 mm. long; petals yellow, sometimes with red spots, sub-orbicular, 3-7 mm. long; capsules clavate, 3-9 mm. long, 1.5 mm. thick; seeds light brown, obovoid, 0.7-1.0 mm. long.

Reported from bare clay hills about St. George, Utah and to the east and south; to be expected in adjacent Nevada.

7. GAYOPHYTUM A. Juss.


GAYOPHYTUM A. Juss. Ann. Sci. Nat. I, 25: 18. pl. 4. 1832.

Slender caulescent annuals; leaves alternate, entire, linear and sessile, or lowest may be opposite and linear-oblongate and short-petioled; flowers in upper axils; hypanthium not prolonged beyond the ovary; sepals 4, usually reflexed in anthesis; petals 4, small, rhomboid-spatulate to -obovate, white, frequently drying pink or red; stamens 8, the alternate set much reduced and usually sterile; stigma capitate; capsule 2-celled, 4-valved, linear or clavate; seeds many, in a single row in each cell, not comose. (Gay, author of Flora of Chile, and Greek word for plant).

A genus of 9 species of the temperate regions of western North America and southern South America; 8 occurring in Nevada.

KEY TO SPECIES

1. Capsule torulose, pedicelled; plants freely branched from the base, repeatedly dichotomous; the upper leaves bract-like.



2. Seeds glabrous.

3. Petals 0.5-1.5 mm. long.

4. The petals 0.5 mm. long; capsule 2-5 mm. long, shorter than the deflexed pedicel; plants quite glabrous.

1. G. ramosissimum.

4. The petals 1-1.5 mm. long; capsule 5-12 mm. long, exceeding the pedicel. 2. G. Nuttallii.

3. Petals 2-4 mm. long. 3. G. diffusum.

2. Seeds appressed-canescant.

5. Petals 1-2 mm. long. 4. G. lasiospermum.

5. Petals 3-4 mm. long. 5. G. eriospermum.

1. Capsule not torulose, subsessile; plants branched mostly at the base, not so much above; upper leaves quite well developed.

6. Seeds vertically placed in a very narrow capsule.

7. The seeds glabrous. 6. G. racemosum.

7. The seeds appressed-canescant. 7. G. Helleri.

6. Seeds obliquely placed in a slightly broader capsule.

8. G. humile.


1. GAYOPHYTUM RAMOSISSIMUM Torr. & Gray, Fl. No. Amer. 1: 513.

1840.

G. ramosissimum var. deflexum Hook. London Journ. Bot.

6: 224. 1847.

Diffusely branched, mostly above the base, with the ultimate branches filiform, quite glabrous sometimes slightly strigulose about the flowers; plant 2-5 dm. tall; leaves lance-linear, 1-3 cm. long, short-petioled, gradually reduced



up the stem; pedicels capillary, 3-5 mm. long, mostly spreading-deflexed; flowers minute; sepals erect, 0.5 mm. long; petals 0.5 mm. long; stigma globose; capsules plump, 2-5 mm. long; seeds glabrous, 0.6 mm. long.

Dry slopes and ridges, Upper Sonoran and Transition Zones, eastern Washington and Oregon to northeastern California, Arizona and Rocky Mts. For Nevada, seen from Washoe, Humboldt, Ormsby, Douglas, Lander, Eureka, Esmeralda, Nye, Elko and White Pine Counties.

2. GAYOPHYTUM NUTTALLII Torr. & Gray, Fl. No. Amer. 1: 514. 1840.


G. ramosissimum var. strictipes Hook. London Journ. Bot.

6: 224. 1847.

With habit and stature of the preceding species, usually more obviously strigulose in the upper parts; pedicels 1-4 mm. long, erect; sepals 1-1.5 mm. long; petals reddish, at least in age, 1-1.5 mm. long; capsules 5-12 mm. long, erect, usually exceeding the pedicels; seeds glabrous, 1-1.5 mm. long.

Dry slopes and ridges, Upper Sonoran and Transition Zones, Washington to Baja California, Dakota, and New Mexico; South America. Nevada plants seen from Washoe, Ormsby, Douglas, Mineral, Esmeralda, Clark, Lander, Elko and White Pine Counties.

var. INTERMEDIUM (Rydb.) Munz, Amer. Journ. Bot. 19: 772. 1932.



G. intermedium Rydb., Bull. Torrey Bot. Club 31: 569. 1904.

Puberulence appressed; pedicels and fruit spreading or deflexed.

Transition Zone, eastern Washington and Oregon to Rocky Mts. For Nevada seen from Washoe, Humboldt, Clark, Elko and Lincoln Counties.

var. ABRAMSII Munz, Amer. Journ. Bot. 19: 772. 1932.

Puberulence short and spreading; pedicels and capsules mostly erect.

Transition Zone, eastern Washington to California and Montana. Nevada plants seen from Washoe County.


3. GAYOPHYTUM DIFFUSUM Torr. & Gray, Fl. No. Amer. 1: 513. 1840.

With general habit and stature of G. Nuttallii, appressed-puberulent in upper parts; pedicels 2-8 mm. long, erect or divaricate; sepals 2-3 mm. long; petals white to pink, 2-4 mm. long; stigma clavate, capitate; capsules 5-12 mm. long, divaricate; seeds glabrous, 1-1.25 mm. long.

Occasional on dry slopes and benches, Transition Zone, Washington to southern California, Montana and Wyoming. Nevada plants seen from Washoe, Douglas and Ormsby Counties.

4. GAYOPHYTUM LASIOSPERMUM Greene, Pittonia 2: 164. 1891.

Plant 2-5 dm. high, branching freely above, with appressed



puberulence in upper parts; leaves linear to lance-linear, 1-3 cm. long, entire; pedicels 3-6 mm. long, divaricate to spreading; sepals 1 mm. long; petals white, turning to rose, 1-2 mm. long; stigma capitate; capsule 4-8 mm. long, torulose; seeds strigose-canescens, 1 mm. long.

Dry places, Transition Zone, Washington to California and Montana. Nevada plants seen from Ormsby, Mineral, Nye, Clark, Lander, and Elko Counties.

5. GAYOPHYTUM ERIOSPERMUM Coville, Contr. U. S. Nat. Herb. 4: 103. 1893.

Like G. lasiospermum in stature, habit, and pubescence of seeds; pedicels 4-8 mm. long; sepals 3 mm. long; petals 3-5 mm. long.

Rare, dry places, Transition Zone, Oregon to California and Idaho. Nevada plants seen from Elko County.

6. GAYOPHYTUM RACEMOSUM Torr. & Gray, Fl. No. Amer. 1: 514. 1840.

Plants low, 1-2 dm. high, subsimple to repeatedly branched from the base, the ultimate branches leafy and relatively simple, strigulose or subglabrous; leaves linear to linear-oblongate, 1-3 cm. long; pedicels from almost none to 2 mm. long, erect; sepals 0.5 mm. long; petals white, turning red, scarcely 1 mm. long; capsule subterete, narrowly linear, not torulose, erect, 6-14 mm. long; seeds erect, glabrous, 1 mm. long.

Dry slopes and flats, Transition Zone, and above; Washington to southern California, Arizona and Rocky Mts. Nevada material seen from Washoe, Humboldt, Eureka, and Elko Counties.

var. CAESIUM (Torr. & Gray) Munz, Amer. Journ. Bot. 19:

776. 1932.

G. caesium Torr. & Gray, Fl. No. Amer. 1: 514. 1840.

Whole plant, or only the upper portion, with minute short spreading hairs.

Not common, Transition Zone, Washington to central California and Nevada, from which state material has been seen for Washoe, Ormsby, Humboldt, Elko and White Pine Counties.

7. GAYOPHYTUM HELLERI Rydb. Bull. Torrey Bot. Club 40: 65. 1913.

With habit, size and aspect of G. racemosum, the puberulence of short spreading hairs; seeds appressed-canescant.

Occasional in dry places, Transition Zone, Washington to southern California and Idaho. Nevada plants seen from Washoe and Ormsby Counties.

var. GLABRUM Munz, Amer. Journ. Bot. 19: 777. 1932.

Plant quite glabrous.

Washington to central California and Colorado. Seen from Washoe and Elko Counties, Nevada.

8. GAYOPHYTUM HUMILE Juss. var. HIRTELLA Munz, Amer. Journ.

Bot. 19: 778. 1932.

Low, 5-15 cm. tall, branched from base, puberulent with short spreading hairs; the branches relatively simple; leaves linear to lance-linear, 1-3 cm. long, entire, on short petioles, upper leaves somewhat reduced, but quite well developed; sepals 1 mm. long; petals white, 1 mm. long; capsule flattened, non-torulose, erect, 10-15 mm. long; seeds obliquely placed in capsules, 0.6 mm. long.

Rare and local, Transition Zone, eastern Central California, eastern Oregon and adjacent Nevada, where it has been reported from Washoe and Ormsby Counties.

♂. GAURA L.

GAURA L. Sp. Pl. 347. 1753.

Cauliscent herbs, annual to perennial; leaves alternate; flowers white or pink, irregular, in terminal racemes or spikes; hypanthium narrow and short; sepals 4, deciduous; petals 4, clawed; stamens usually 8, with or without scale-like appendage at base of each filament; ovary 4-celled, with single ovule in each cell; stigma 4-lobed, with cup-like border at base, or discoid and entire; capsule nut-like, obovoid, nearly or quite indehiscent, 1- to 4-seeded. (Greek, proud, some species being showy).

A genus of 16 species of North and South America, 2 occurring in Nevada.

KEY TO SPECIES

1. Plants 5-20 dm. tall, biennial; leaves 1-3 cm. wide, 5-10 cm.

long; anthers oval, attached near the middle.

1. G. parviflora.

1. Plants 1-4 dm. tall, perennial; leaves 0.3-1 cm. wide, 1-4 cm.

long; anthers linear, attached near the base. . 2. G. coccinea.

1. GAURA PARVIFLORA Dougl. ex Hook. Fl. Bor. Amer. 1: 208. 1834.


Biennial or winter annual, erect, 5-20 dm. tall, simple or with few branches, silky-pilose, with long spreading hairs on stems, veins and leaf-margins, and also with minute glandular pubescence; leaves ovate-lanceolate, almost sessile, repand-dentate to subentire, 3-10 cm. long, 1-3 cm. wide, finely pubescent on general surface; inflorescence spicate, 1-3 dm. long; flowers numerous; hypanthium puberulent, 2-3 mm. long; sepals glabrous to puberulent, 2-3 mm. long; petals 2-4 mm. long, reddish; fruit sessile, 6-10 mm. long, somewhat fusiform, glabrous; seeds 1-2, brown.

Disturbed and waste places, Upper Sonoran and Transition Zones; eastern Washington to Mississippi Valley and Mexico. Reported from Battle Mountain, Lander Co., Nevada.

2. GAURA COCCINEA NUTT. in Fraser's Cat., 1813 as nomen nudum;

ex Pursh, Fl. Sept. Amer. 2: 733. 1816.

Perennial herb; stems several to many, simple or branched, ascending, canescent-strigose, mostly 1-3 dm. high; leaves numerous, sessile, oblong-lanceolate to linear, entire to



repand-denticulate, 1-3 cm. long, 3-10 mm. wide; spikes short, 1-2 dm. long; hypanthium 6-10 mm. long; sepals 6-9 mm. long; petals pink or red, turning scarlet, 5-8 mm. long; stamens almost as long as petals; fruits canescent, 5-7 mm. long, 2.5-3 mm. thick, the body proper 4-angled, abruptly constricted into a stout base.

Dry slopes, Upper Sonoran Zone, Alberta and Manitoba to southeastern California, Arizona and Texas. Fairly common in Clark County, Nevada.

var. GLABRA (Lehm.) Torr. & Gray, Fl. No. Amer. 1: 518.

1840.

Plant nearly or quite glabrous; leaves more wavy; hypanthium strigulose.

With much the range of the species. Clark Co., Nevada.

9. CIRCAEA L.

CIRCAEA L. Sp. Pl. 9. 1753.

Low slender perennial herbs with subterranean rootstocks with tuber-like enlargements; leaves opposite, thin, petioled; flowers small, in racemes paniculately disposed; hypanthium short, deciduous and with a ring-like disk within; sepals 2, reflexed; petals 2, white, notched; stamens 2, alternate with the petals; ovary 1- or 2-celled, each cell 1-ovuled; fruit nut-like, 1- or 2-seeded, obovoid, indehiscent, usually with hooked hairs. (Named for Circe, the enchantress).

Several species, from cooler places in the northern hemisphere; one occurring in Nevada.

1. *CIRCAEA PACIFICA* Aschers. & Magnus. Bot. Zeit. 29: 392. 1871.

Erect, 2-4 dm. tall; leaf-blades ovate, usually rounded at the base, entire or minutely denticulate, 2-6 cm. long, on petioles 2-3 cm. long; pedicels 3-4 mm. long, reflexed in fruit; calyx and corolla about 1 mm. long; capsule narrowly obovoid, 1-celled, 2 mm. long, covered with soft, hooked hairs.

Cold moist woods of the Transition Zone, British Columbia to southern California and the Rocky Mts. Nevada collections seen from Washoe, Douglas, and Elko Counties.

INDEX

	PAGE	EPILOBIUM	PAGE
Anogra		LATIFOLIUM	5,9
longiflora	40	MINUTUM	6,12
BOISDUVALIA	2,20	NEVADENSE	5,10
DENSIFLORA	21	OBCORDATUM	5,9
var. glabella	21,22	occidentale	19
STRICTA	21,23	OREGONENSE	7,16
Chamaenerion		ovatifolium	14
angustifolium	8	Palmeri	20
latifolium	9	PANICULATUM	6,11
Chylismia		forma ADENOCADON	11
brevipes	56	forma LAEVICAULE	12
clavaeformis	59	forma SUBLATUM	11
heterochroma	60	forma TRACYI	11
hirta	57	platyphyllum	16
multijuga	57	SAXIMONTANUM	7,14
Parryi	61	spicatum	8
pterosperma	55	GAURA	2,68
scapoidea	58	COCCINEA	69
venosa	57	var. GLABRA	70
CIRCAEA	1,70	PARVIFLORA	69
PACIFICA	71	GAYOPHYTUM	2,62
CLARKIA	2,24	caesium	67
RHOMBOIDEA	24	DIFFUSUM	63,65
EPILOBIUM	2,4	ERIOSPERMUM	63,66
ADENOCADON	8,19	HELLERI	63,67
var. OCCIDENTALE	19	var. GLABRUM	67
var. PERPLEXANS	8,20	HUMILE	63
affine		var. HIRTELLA	67
var. fastigiatum	16	intermedium	65
ALPINUM	7,16,18	LASIOSPERMUM	63,65,66
anagallidifolium	16	NUTTALLII	64,65
ANGUSTIFOLIUM	5,8	var. ABRAMSII	65
BREVISTYLUM	6,13	var. INTERMEDIUM	64
var. TENUE	14	RACEMOSUM	63,66,67
var. URSINUM	13	var. CAESIUM	67
CALIFORNICUM	8,20	RAMOSISSIMUM	63
CLAVATUM	7,17	var. deflexum	63
delicatum		var. strictipes	64
var. tenue	14	strictum	23
Drummondii	14	GODETIA	3,25
EXALTATUM	6,12	AMOENA	26
GLABERRIMUM	7,15	latifolia	24
var. FASTIGIATUM	15	Lavauxia	
glandulosum		flava	43
var. adenocaulon	19	Howardi	44
HALLEANUM	6,14	lobata	41
HORNEMANII	8,17	Onagra	
LACTIFLORUM	8,18	strigosa	34

OENOTHEIRA	PAGE
ALYSSOIDES	31
var. VILLOSA	50
amoena	26
ANDINA	32,52
BOOTHII	31,52
BRACHYCARPA	27
var. WRIGHTII	44
BREVIPIES	32,56
var. pallidula	57
var. parviflora	58
CAESPITOSA	28
var. CRINITA	40
var. LONGIFLORA	40
var. MARGINATA	38,40
var. MONTANA	39
var. PURPUREA	40
CALIFORNICA	29,38
CAVERNAE	28,41
CHAMAENERIOIDES	30,48
CLAVAEFORMIS	33,59
var. AURANTIACA	60
var. cruciformis	60
var. PURPURASCENS	60
CONTORTA	32,53
var. FLEXUOSA	54
var. PUBENS	54
DECORTICANS	30,31
var. CONDENSATA	51
var. DESERTORUM	30,51
DELTOIDES	29,36
var. ambigua	36
var. DECUMBENS	36
forma GLABRATA	37
var. PIPERI	36
densiflora	21
DENTATA	32
var. JOHNSTONII	55
FLAVA	27,43
glabella	22
HETERANTHA	29,46
var. taraxacifolia	46
HETEROCHROMA	33,60
var. MEGALANTHA	61
HOOKERI	28
var. ANGUSTIFOLIA	34
LAVANDULAEFOLIA	30
var. GLANDULOSA	44
LONGISSIMA	28,35
marginata	38
var. purpurea	40
MINOR	30,49

OENOTHEIRA	PAGE
montana	39
MULTIJUGA	33,57
var. PARVIFLORA	58
NEVADENSIS	31,49
PALLIDA	29,37
PALLIDULA	32,57
PALMERI	29,45
PARRYI	33,61
PRIMIVERIS	28,41
PTEROSPERMA	32,55
REFRACTA	30,47
RYDBERGII	28,34
SCAPOIDEA	33,58
var. aurantiaca	60
var. purpurascens	60
strigosa	34
var. pubens	54
subacaulis	46
TANACETIFOLIA	29,47
tenuissima	61
Wrightii	44
XYLOCARPA	28,42
Onagra	
strigosa	34
Pachylophus	
canescens	40
crinitus	40
Sphaerostigma	
andinum	53
Boothii	52
chamaenerioides	48
contortum	53
flexuosum	54
macrophyllum	50
minus	49
nevadense	50
orthocarpum	54
refractum	47
senex	52
tortum	49
tortuosum	50
utahense	50
Taraxia	
Palmeri	45
tanacetifolia	47
ZAUSCHNERIA	2,3
argentea	4
californica	
var. latifolia	3
subsp. latifolia	3
LATIFOLIA	3

[Faint, illegible text, likely bleed-through from the reverse side of the page]

[Faint, illegible text, likely bleed-through from the reverse side of the page]

305 1
17-①